

5000


09600660



PCT

 D INTELLECTUAL PROPERTY ORGANIZATION
 International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : C07D 9/38, 213/54, 211/34, A61K 31/662, 31/44, 31/445	A1	(11) International Publication Number: WO 00/66550 (43) International Publication Date: 9 November 2000 (09.11.00)
(21) International Application Number: PCT/SE00/00846 (22) International Filing Date: 3 May 2000 (03.05.00) (30) Priority Data: 9901572-9 3 May 1999 (03.05.99) SE (71) Applicant (for all designated States except US): AS- TRAZENECA AB [SE/SE]; S-151 85 Södertälje (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): LINSCHOTEN, Marcel [NL/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE). POLLA, Magnus [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE). (74) Agent: ASTRAZENECA AB; Global Intellectual Property, Patents, S-151 85 Södertälje (SE).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: NEW COMPOUNDS <div style="text-align: center;">  (1) </div> (57) Abstract <p>The present invention relates to compounds of Formula (I), and pharmaceutically acceptable salts or solvates thereof, or solvates of such salts, which compounds inhibit carboxypeptidase U and thus can be used in the prevention and treatment of diseases associated with carboxypeptidase U. In further aspects, the invention relates to compounds of the invention for use in therapy; to processes for preparation of such new compounds; to pharmaceutical compositions containing at least one compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, as active ingredient; and to the use of the active compounds in the manufacture of medicaments for the medical use indicated above.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

NEW COMPOUNDS

FIELD OF THE INVENTION

5 The present invention relates to novel compounds, and pharmaceutically acceptable salts thereof, which inhibit basic carboxypeptidases, more specifically carboxypeptidase U, and thus can be used in the prevention and treatment of diseases wherein inhibition of carboxypeptidase U is beneficial. In further aspects, the invention relates to compounds of the invention for use in therapy; to processes for preparation of such new compounds; to
10 pharmaceutical compositions containing at least one compound of the invention, or a pharmaceutically acceptable salt thereof, as active ingredient; and to the use of the active compounds in the manufacture of medicaments for the medical use indicated above.

BACKGROUND OF THE INVENTION

15 Fibrinolysis is the result of a series of enzymatic reactions resulting in the degradation of fibrin by plasmin. The activation of plasminogen is the central process in fibrinolysis. The cleavage of plasminogen to produce plasmin is accomplished by the plasminogen activators, tissue-type plasminogen activator (t-PA) or urokinase-type plasminogen
20 activator (u-PA). Initial plasmin degradation of fibrin generates carboxy-terminal lysine residues that serves as high affinity binding sites for plasminogen. Since plasminogen bound to fibrin is much more readily activated to plasmin than free plasminogen this mechanism provides a positive feedback regulation of fibrinolysis.

25 One of the endogenous inhibitors to fibrinolysis is carboxypeptidase U (CPU). CPU is also known as plasma carboxypeptidase B, active thrombin activatable fibrinolysis inhibitor (TAFIa), carboxypeptidase R and inducible carboxypeptidase activity. CPU is formed during coagulation and fibrinolysis from its precursor proCPU by the action of proteolytic enzymes *e.g.* thrombin, thrombin-thrombomodulin complex or plasmin. CPU cleaves basic
30 amino acids at the carboxy-terminal of fibrin fragments. The loss of carboxy-terminal lysines and thereby of lysine binding sites for plasminogen then serves to inhibit fibrinolysis.

By inhibiting the loss of lysine binding sites for plasminogen and thus increase the rate of plasmin formation, effective inhibitors of carboxypeptidase U would be expected to facilitate fibrinolysis.


2-mercaptomethyl-3-guanidinoethylthiopropionic acid is reported as a carboxypeptidase N inhibitor. More recently, this compound has been shown to inhibit CPU, Hendriks, D. *et al.*, Biochimica et Biophysica Acta, 1034 (1990) 86-92.

Guanidinoethylmercaptosuccinic acid is reported as a carboxypeptidase N inhibitor. More recently, this compound has been shown to inhibit CPU, Eaton, D. L., *et al.*, The Journal of Biological Chemistry, 266 (1991) 21833-21838.

DISCLOSURE OF THE INVENTION

It has surprisingly been found that compounds of the Formula I are particularly effective as inhibitors of carboxypeptidase U and thereby useful as medicaments for the treatment or prophylaxis of conditions wherein inhibition of carboxypeptidase U is beneficial.

In one aspect, the invention thus relates to compounds of the general Formula I,



or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt, wherein

R₁ represents,

C₁-C₆ alkyl, substituted with one or more basic groups such as amino, amidino and/or guanidino;

cycloalkyl, substituted with one or more basic groups such as amino, amidino and/or guanidino;

heterocyclyl, containing at least one nitrogen atom;

heterocyclyl, containing at least one hetero atom selected from S or O,

5 and substituted with one or more basic groups such as amino, amidino and/or guanidino;

or aryl, substituted with one or more basic groups such as amino, amidino and/or guanidino,

R_2 represents H, acyl, acylamino, alkyl, alkylcarbamoyl, alkylthio, alkoxy, aroyl, aroylamino, aryloxy, arylthio, amidino, amino, aryl, carbamoyl, carboxy, cyano, cycloalkyl, formyl, guanidino, halogen, heterocyclyl, hydroxy, oxo, nitro, thiol, Z_2N -CO-O-, ZO-CO-NZ- or Z_2N -CO-NZ- group,

R_3 represents $COOR_5$, $SO(OR_5)$, SO_3R_5 , $P=O(OR_5)_2$, $B(OR_5)_2$, $P=OR_5(OR_5)$, or tetrazole, or any carboxylic acid isostere,

15

R_4 represents a $\begin{array}{c} \text{O}-R_5 \\ | \\ -P- \\ || \\ \text{O} \end{array} R_6$ -group, or a $\begin{array}{c} \text{O} \\ || \\ \text{C} \\ | \\ \text{N}-\text{OH} \\ | \\ R_7 \end{array}$ -group, or a $\begin{array}{c} \text{O} \\ || \\ \text{C} \\ | \\ \text{O}-R_5 \end{array}$ -group,

R_5 represents H, C_1 - C_6 alkyl or aryl,

R_6 represents C_1 - C_6 alkyl, aryl, cycloalkyl, heterocyclyl, or an optionally N-substituted H_2N -C(Z)-CONH-C(Z)- or H_2N -C(Z)- group,

20 R_7 represents H or C_1 - C_6 alkyl,

X represents O, S, SO, SO_2 , $C(Z)_2$, N(Z), NR_7SO_2 , SO_2NR_7 , NR_7CO or $CONR_7$,

Y represents O, N(Z), S, $C(Z)_2$, or a single bond,

Z represents independently H, C_1 - C_6 alkyl, aryl, cycloalkyl or heterocyclyl,

with the proviso that when X represents O, S, SO, SO_2 , N(Z), NR_7SO_2 , SO_2NR_7 , or

25 NR_7CO then Y represents $C(Z)_2$ or a single bond.

Preferred compounds according to the present invention are those of Formula I, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt,

wherein

R₁ represents,

C₁-C₆ alkyl, substituted with one or more basic groups such as amino, amidino and/or guanidino;

5 cycloalkyl, substituted with one or more basic groups such as amino, amidino and/or guanidino;

heterocyclyl, containing at least one nitrogen atom;

heterocyclyl, containing at least one hetero atom selected from S or O,

and substituted with one or more basic groups such as amino, amidino and/or

10 guanidino;




or aryl, substituted with one or more basic groups such as amino, amidino and/or guanidino,

R₂ represents H, acyl, acylamino, alkyl, alkylcarbamoyl, alkylthio, alkoxy, aroyl,

aroylamino, aryloxy, arylthio, amidino, amino, aryl, carbamoyl, carboxy, cyano,

15 cycloalkyl, formyl, guanidino, halogen, heterocyclyl, hydroxy, oxo, nitro, thiol, Z₂N-CO-O-, ZO-CO-NZ- or Z₂N-CO-NZ- group,

R₃ represents COOR₅,

R₄ represents a  -group, or a  -group, or a  -group,

20 R₅ represents H, C₁-C₆ alkyl or aryl,

R₆ represents C₁-C₆ alkyl, aryl, cycloalkyl, heterocyclyl, or an optionally N-substituted

H₂N-C(Z)-CONH-C(Z)- or H₂N-C(Z)- group,

R₇ represents H or C₁-C₆ alkyl,

X represents C(Z)₂,

25 Y represents O, N(Z), S, C(Z)₂, or a single bond,

Z represents independently H, C₁-C₆ alkyl, aryl, cycloalkyl or heterocyclyl.

More preferred compounds according to the present invention are those of Formula I, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt, wherein

R₁ represents,

- 5 cycloalkyl, substituted with one or more basic groups such as amino, amidino and/or guanidino;
 heterocyclyl, containing at least one nitrogen atom;
 heterocyclyl, containing at least one hetero atom selected from S or O,
 and substituted with one or more basic groups such as amino, amidino and/or
 10 guanidino;

R₂ represents H, C₁-C₃ alkyl, amino, halogen or hydroxy,

R₃ represents COOR₅,

R₄ represents a $\begin{array}{c} \text{O}-\text{R}_5 \\ | \\ -\text{P}-\text{R}_6 \\ || \\ \text{O} \end{array}$ -group,

R₅ represents H, C₁-C₆ alkyl or aryl,

- 15 R₆ represents C₁-C₆ alkyl, aryl, cycloalkyl, heterocyclyl, or an optionally N-substituted
 H₂N-C(Z)-CONH-C(Z)- or H₂N-C(Z)- group,

X represents C(Z)₂,

Y represents O or C(Z)₂,

Z represents independently H or C₁-C₆ alkyl.

20

Other more preferred compounds according to the present invention are those of Formula I, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt, wherein


R₁ represents,

- 25 cycloalkyl, substituted with one or more basic groups such as amino, amidino and/or guanidino;
 heterocyclyl, containing at least one nitrogen atom;

heterocyclyl, containing at least one hetero atom selected from S or O,
and substituted with one or more basic groups such as amino, amidino and/or
guanidino;

R₂ represents H, C₁-C₃ alkyl, amino, halogen or hydroxy,

5 R₃ represents COOR₅,

R₄ represents a  -group,

R₅ represents H, C₁-C₆ alkyl or aryl,

R₇ represents H or C₁-C₆ alkyl,

X represents C(Z)₂,

10 Y represents C(Z)₂ or a single bond,

Z represents independently H or C₁-C₆ alkyl.

Yet other more preferred compounds according to the present invention are those of
Formula I, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a
15 salt,

wherein

R₁ represents,


cycloalkyl, substituted with one or more basic groups such as amino, amidino and/or
guanidino;

20 heterocyclyl, containing at least one nitrogen atom;

heterocyclyl, containing at least one hetero atom selected from S or O,
and substituted with one or more basic groups such as amino, amidino and/or
guanidino;

R₂ represents H, C₁-C₃ alkyl, amino, halogen or hydroxy,

25 R₃ represents COOR₅,

R₄ represents a -group,

R₅ represents H, C₁-C₆ alkyl or aryl,

X represents C(Z)₂,

Y represents C(Z)₂ or a single bond,

5 Z represents independently H or C₁-C₆ alkyl.

Even more preferred compounds according to the present invention are those of Formula I, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt, wherein


10 R₁ represents,

cycloalkyl, substituted with one or more basic groups such as amino, amidino and/or guanidino;

heterocyclyl, containing at least one nitrogen atom;

R₂ represents H, F, C₁ alkyl,

15 R₃ represents COOR₅,

R₄ represents a -group,

R₅ represents H, C₁-C₆ alkyl or aryl,

R₆ represents C₁-C₆ alkyl, aryl, cycloalkyl, heterocyclyl, or an optionally N-substituted

H₂N-C(Z)-CONH-C(Z)- or H₂N-C(Z)- group,

20 X represents C(Z)₂,

Y represents O or C(Z)₂,

Z represents independently H or C₁-C₆ alkyl.

Other even more preferred compounds according to the present invention are those of
25 Formula I, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt,

wherein


R₁ represents,

cycloalkyl, substituted with one or more basic groups such as amino, amidino and/or guanidino;

5 heterocyclyl, containing at least one nitrogen atom;

R₂ represents H, F, C₁ alkyl,

R₃ represents COOR₅,

R₄ represents a  -group,

R₅ represents H, C₁-C₆ alkyl or aryl,

10 R₇ represents H or C₁-C₆ alkyl,

X represents C(Z)₂,

Y represents C(Z)₂ or a single bond,

Z represents independently H or C₁-C₆ alkyl.

15 Yet other even more preferred compounds according to the present invention are those of Formula I, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt,

wherein

R₁ represents,


20 cycloalkyl, substituted with one or more basic groups such as amino, amidino and/or guanidino;

heterocyclyl, containing at least one nitrogen atom;

R₂ represents H, F, C₁ alkyl,

R₃ represents COOR₅,

25

R₄ represents a  -group,

R₅ represents H, C₁-C₆ alkyl or aryl,

X represents C(Z)₂,

Y represents C(Z)₂ or a single bond,

Z represents independently H or C₁-C₆ alkyl.


5

Most preferred compounds according to the present invention are those of Formula I, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt, wherein

R₁ represents pyridyl or piperidinyl,

10 R₂ represents H,

R₃ represents COOR₅,

R₄ represents a  -group,

R₅ represents H,

R₆ represents C₁-C₆ alkyl or an optionally N-substituted H₂N-C(Z)-CONH-C(Z)- or H₂N-

15 C(Z)- group,

X represents CHZ,

Y represents CHZ,


Z represents independently H or C₁-C₆ alkyl.

20 Other most preferred compounds according to the present invention are those of Formula I, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt, wherein

R₁ represents pyridyl or piperidinyl,

R₂ represents H,

25 R₃ represents COOR₅,

R₄ represents a  -group,

R₅ represents H,

R₇ represents H,

X represents CHZ,

5 Y represents CHZ or a single bond,

Z represents independently H or C₁-C₆ alkyl.

Yet other most preferred compounds according to the present invention are those of
Formula I, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a
10 salt,

wherein

R₁ represents pyridyl or piperidiny1,

R₂ represents H,

R₃ represents COOR₅,

R₄ represents a  -group,

15 R₅ represents H,

X represents CHZ,

Y represents CHZ or a single bond,

Z represents independently H or C₁-C₆ alkyl.

20 The following definitions shall apply throughout the specification and the appended
claims:

The term "basic group" denotes a basic group, wherein the conjugate acid of said basic
25 group has a pK_a of from about -5 to about 25, preferably of from 1 to 15.

The term "carboxylic acid isostere" denotes an acidic group having a pK_a of from about -5
to about 25, preferably of from 1 to 15.

The term "C₁-C₆ alkyl" denotes a straight or branched, saturated or unsaturated, substituted or unsubstituted alkyl group having 1 to 6 carbon atoms in the chain wherein the alkyl group may optionally be interrupted by one or more heteroatoms selected from O, N or S. Examples of said alkyl include, but is not limited to, methyl, ethyl, ethenyl, ethynyl, n-propyl, iso-propyl, propenyl, iso-propenyl, propynyl, n-butyl, iso-butyl, sec-butyl, t-butyl, butenyl, iso-butenyl, butynyl and straight- and branched-chain pentyl and hexyl.

The term "C₁-C₃ alkyl" denotes a straight or branched, saturated or unsaturated, substituted or unsubstituted alkyl group having 1 to 3 carbon atoms in the chain wherein the alkyl group may optionally be interrupted by one or more heteroatoms selected from O, N or S. Examples of said alkyl include, but is not limited to, methyl, ethyl, ethenyl, ethynyl, n-propyl, iso-propyl, propenyl, iso-propenyl, propynyl.

The term "C₁ alkyl" denotes a substituted or unsubstituted alkyl group having 1 carbon atom. An example of said alkyl include, but is not limited to, methyl,

The term "C₁-C₆ alkoxy" denotes an alkyl-O-group, wherein C₁-C₆ alkyl is as defined above.

The term "C₁-C₃ alkoxy" denotes an alkyl-O-group, wherein C₁-C₃ alkyl is as defined above.

The term "heterocyclyl" denotes a substituted or unsubstituted, 4- to 10- membered monocyclic or polycyclic ring system in which one or more of the atoms in the ring or rings is an element other than carbon, for example nitrogen, oxygen or sulfur, especially 4-, 5- or 6-membered aromatic or aliphatic heterocyclic groups, and includes, but is not limited to azetidine, furan, thiophene, pyrrole, pyrrolidine, dioxolane, oxathiolane, oxazoline, oxazole, thiazole, imidazole, imidazoline, imidazolidine, pyrazole, pyrazoline, pyrazolidine, isoxazole, isothiazole, oxadiazole, furazan, triazole, thiadiazole, pyran, pyridine, piperidine, dioxane, morpholine, dithiane, oxathiane, thiomorpholine, pyridazine, pyrimidine, pyrazine, piperazine, triazine, thiadiazine, dithiazine, azaindole, azaindoline,

indole, indoline, naphthyridine groups, and shall be understood to include all isomers of the above identified groups. The term "azetidiny" shall for example be understood to include the 2-, and 3-isomers and the terms "pyridyl" and "piperidiny" shall for example be understood to include the 2-, 3-, and 4-isomers.

5

The term "cycloalkyl" denotes a saturated or unsaturated, substituted or unsubstituted, non-aromatic ring composed of 3, 4, 5, 6 or 7 carbon atoms, and includes, but is not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclopentadienyl, cyclohexadienyl and cyclo-

10

heptadienyl groups,

The term "halogen" includes fluoro, chloro, bromo and iodo groups.

The term "aryl" denotes a substituted or unsubstituted C₆-C₁₄ aromatic hydrocarbon and includes, but is not limited to, phenyl, naphthyl, indenyl, anthracenyl, fenantrenyl, and fluorenyl.

15

The term "aryloxy" denotes an aryl-O-group, wherein aryl is as defined above.

The term "acyl" denotes an alkyl-CO-group, wherein alkyl is as defined above.

20

The term "aroyl" denotes an aryl-CO-group, wherein aryl is as defined above.

The term "alkylthio" denotes an alkyl-S-group, wherein alkyl is as defined above.

25

The term "arylthio" denotes an aryl-S-group, wherein aryl is as defined above.

The term "aroylamino" denotes an aroyl-N(Z)-group, wherein aroyl and Z is as defined above.

30

The term "acylamino" denotes an acyl-N(Z)-group, wherein acyl and Z is as defined above.

The term "carbamoyl" denotes a $\text{H}_2\text{N-CO}$ -group.

The term "alkylcarbamoyl" denotes a $\text{Z}_2\text{N-CO}$ -group wherein Z is as defined above.

5

The term "substituted" denotes an " C_1 alkyl", " $\text{C}_1\text{-C}_3$ alkyl", " $\text{C}_1\text{-C}_6$ alkyl", "cycloalkyl", "heterocyclyl", "aryl", $\text{H}_2\text{N-C(Z)-CONH-C(Z)-}$ or a $\text{H}_2\text{N-C(Z)-}$ group as defined above which is substituted by one or more acyl, acylamino, alkyl, alkylcarbamoyl, alkylthio, alkoxy, aroyl, aroylamino, aryloxy, arylthio, amidino, amino, aryl, carbamoyl, carboxy, cyano, cycloalkyl, formyl, guanidino, halogen, heterocyclyl, hydroxy, oxo, nitro, thiol, thio, $\text{Z}_2\text{N-CO-O-}$, ZO-CO-NZ- , or $\text{Z}_2\text{N-CO-NZ-}$ groups.

10

Both the pure enantiomers, racemic mixtures and unequal mixtures of two enantiomers are within the scope of the present invention. It should also be understood that all the diastereomeric forms possible are within the scope of the invention. Also included in the invention are derivatives of the compounds of the Formula I which have the biological function of the compounds of Formula I, such as prodrugs.

15

Depending on the process conditions the compounds of Formula I are obtained either in neutral or salt form or as a solvate, e.g. a hydrate, and are all within the scope of the present invention.

20

Preparation


The present invention also provides the process A-C for the manufacture of compounds with the general Formula I.

25

Process A

Process A for manufacture of compounds with the general Formula I, wherein R_1 , R_5 , R_6 , and Z are as defined above and R_2 is H, R_3 is COOR_5 ,

30

R₄ represents a -group,

X is C(Z)₂, Y is C(Z)₂ and comprises the following steps:

a) Compounds of the general Formula II,




5 wherein R₁ and Z is as defined for Formula I and X is C(Z)₂, which are either commercially available or are available using known techniques, can be converted into a compound of the general Formula III,




10 wherein L is a suitable leaving group, such as a chloro, bromo, iodo, triflate or tosyl group, under standard conditions using a suitable reagent, such as PPh₃/CBr₄, TosCl/pyridine or (CF₃SO₂)₂O/TEA).

b) Compounds of the general Formula III can thereafter be reacted with compounds of the general Formula IV,


15



20 which are either commercially available or are available using known techniques, in the presence of a suitable base, such as K₂CO₃ or NaH, under standard conditions to give compounds of the general Formula V,




c) Compounds of the general Formula V can thereafter be converted to compounds of the general Formula VI,



5 by treatment with formaldehyd in the presence of a suitable base, such as Et_2NH , under standard conditions.

However, if Y is $\text{CH}(\text{Z})$ then compounds of the general Formula VI can be prepared by treating compounds of the general Formula VII,


10



with an alkylating agent of the general Formula III,




15 wherein R_1 is as defined for Formula I and L is a suitable leaving group, such as a chloro, bromo, iodo, triflate or tosyl group, in the presence of a suitable base, such as LDA or NaH, under standard conditions to give compounds of the general Formula VIII,




Compounds of the general Formula VIII can thereafter be reacted with an appropriate aldehyde CHO(Z), wherein Z is as defined for Formula I, in the presence of a suitable base, such as KOtBu, LDA or NaH, under standard conditions to give a compound of the general Formula VI.

d) Compounds of the general Formula VI can be further reacted with compounds of the general Formula IX



wherein R_6 is as defined for Formula I, in the presence of a suitable reagent, such as BSA or HMDS, under standard conditions to give compounds of the general Formula I, wherein R_1 , R_5 , R_6 and Z are as defined above, R_2 is H, R_3 is COOR₅,




X is C(Z)₂, and Y is C(Z)₂.

15

Process B


Process B for manufacture of compounds with the general Formula I, wherein R_1 , R_2 , R_5 , R_6 , and Z are as defined above, R_3 is COOR₅, X is C(Z)₂, Y is O, and



20


comprises the following steps:

a) Reacting a compound of the general Formula X,



wherein R_1 , R_2 and Z are as defined in Formula I and X is $C(Z)_2$ in the presence of suitable reagents, such as $TMSCN/ZnI_2$ or $KCN/HOAc$, under standard conditions to give compounds of the general Formula XI,


5



wherein R_1 and R_2 are as defined in Formula I and X is $C(Z)_2$.

b) Compounds of the general Formula XI can thereafter be treated with suitable reagents, such as HCl or $HCl/MeOH$, under standard conditions to give compounds of the general Formula XII,

10




wherein R_1 and R_2 are as defined in Formula I and X is $C(Z)_2$.

15

c) Compounds of the general Formula XII can thereafter be reacted with compounds of the general Formula XIII,

20



wherein R_6 is as defined in the general Formula I, which are either commercially available, well known in the literature, or are available using known techniques, in the presence of suitable coupling reagents such as $DCC/DMAP$, $PyBop/DIPEA$ or $SOCl_2$, under standard

conditions to give compounds of the general Formula I, wherein R_1 , R_2 , R_5 , R_6 and Z are as defined above, R_3 is COOR_5 ,

R_4 represents a $\begin{array}{c} \text{O}-R_5 \\ | \\ -\text{P}-R_6 \\ || \\ \text{O} \end{array}$ -group,


X is $\text{C}(\text{Z})_2$ and Y is O .

5

Process C

Process C for manufacture of compounds with the general Formula I, wherein R_1 and R_2 are as defined above and X and Y is $\text{C}(\text{Z})_2$ or a single bond and R_3 and R_4 are COOR_5 ,
 10 comprises the following steps,


a) reacting a compound of the general Formula XIV,



15 wherein R_2 and R_5 are as defined in Formula I and Y is $\text{C}(\text{Z})_2$ or a single bond, which are either commercially available, well known in the literature, or are available using known techniques, with a compound of the general Formula III,



wherein R_1 is as defined for Formula I, X is $\text{C}(\text{Z})_2$ and L is a suitable leaving group, such
 20 as Cl , Br , I or tosyl, in the presence of a suitable base, such as LDA or NaH under standard conditions, to give a compound of the general Formula XV,




- b) hydrolysing a compound of the general Formula XV, for example by treatment with aqueous NaOH or aqueous TFA under standard conditions to give compounds of the general Formula I, wherein R_1 and R_2 are as defined above and X and Y is $\text{C}(\text{Z})_2$ or a single bond and R_3 and R_4 are COOH .


Process D

10

Process D for manufacture of compounds with the general Formula I, wherein R_1 , R_2 , R_5 , R_7 , X, Y and Z are as defined above, R_3 is COOR_5 and

R_4 represents a  -group, comprises the following steps:

- 15 a) Compounds of the general Formula XV,




can be reacted with compounds of the general Formula XVI,

20



wherein R_7 is as defined in Formula I, in the presence of suitable reagents, such as DCC/DMAP, under standard conditions, to give compounds of the general Formula I, wherein R_1 , R_2 , R_5 , R_7 , X, Y and Z are as defined above, R_3 is COOR_5 and

5

R_4 represents a  -group.

It will be appreciated by those skilled in the art that in the process described above the functional groups of intermediate compounds may need to be protected by protecting groups.

10

Functional groups which it is desirable to protect include hydroxy, amino, mercapto and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl or diarylalkylsilyl (e.g. t-butyldimethylsilyl, t-butyldiphenylsilyl or trimethylsilyl), tetrahydropyranyl and benzyl. Suitable protecting groups for amino, amidino and guanidino include t-butyloxycarbonyl and benzyloxycarbonyl. Suitable protecting groups for mercapto include CO- C_{1-6} alkyl, p-methoxybenzyl and trityl. Suitable protecting groups for carboxylic acid include C_{1-6} alkyl and benzyl esters.

15

Protecting groups may be removed in accordance with techniques which are well known to those skilled in the art and as described hereinafter.

20

Certain protected derivatives of compounds of Formula I, which may be made prior to a final deprotection stage to form compounds of Formula I, are novel.

25

The use of protecting groups is described in 'Protective Groups in Organic Synthesis', 2nd edition, T.W. Greene & P.G.M. Wutz, Wiley-Interscience (1991). The protective group may also be a polymer resin such as Wang resin or a 2-chlorotriptyl chloride resin.

5 It will also be appreciated by those skilled in the art, although such protected derivatives of compounds of Formula I may not possess pharmacological activity as such, they may be administered parenterally or orally and thereafter metabolised in the body to form compounds of the invention which are pharmacologically active. Such derivatives may therefore be described as "prodrugs". All prodrugs of compounds of Formula I are included
10 within the scope of the invention.

It should also be understood that all polymorphs, amorphous forms, anhydrides, hydrates, solvates of the compounds of the present invention are within the scope of the invention.

15 *Pharmaceutical formulations*

In yet a further aspect, the invention relates to pharmaceutical compositions containing at least one compound of the present invention, or a pharmaceutically acceptable salt thereof, as active ingredient.

20 For clinical use, the compounds of the invention are formulated into pharmaceutical formulations for oral, intravenous, subcutaneous, tracheal, bronchial, intranasal, pulmonary, transdermal, buccal, rectal, parenteral or other mode of administration. The pharmaceutical formulation contains a compound of the invention in combination with one
25 or more pharmaceutically acceptable ingredients. The carrier may be in the form of a solid, semi-solid or liquid diluent, or a capsule. These pharmaceutical preparations are a further object of the invention. Usually the amount of active compounds is between 0.1–95% by weight of the preparation.

30 In the preparation of pharmaceutical formulations containing a compound of the present invention the compound selected may be mixed with solid, powdered ingredients, such as lactose, saccharose, sorbitol, mannitol, starch, amylopectin, cellulose derivatives, gelatin,

or another suitable ingredient, as well as with disintegrating agents and lubricating agents such as magnesium stearate, calcium stearate, sodium stearyl fumarate and polyethylene glycol waxes. The mixture may then be processed into granules or pressed into tablets.

5 Soft gelatine capsules may be prepared with capsules containing a mixture of the active compound or compounds of the invention, vegetable oil, fat, or other suitable vehicle for soft gelatine capsules. Hard gelatine capsules may contain granules of the active compound. Hard gelatine capsules may also contain the active compound in combination with solid powdered ingredients such as lactose, saccharose, sorbitol, mannitol, potato
10 starch, corn starch, amylopectin, cellulose derivatives or gelatine.

Dosage units for rectal administration may be prepared (i) in the form of suppositories which contain the active substance mixed with a neutral fat base; (ii) in the form of a gelatine rectal capsule which contains the active substance in a mixture with a vegetable
15 oil, paraffin oil or other suitable vehicle for gelatine rectal capsules; (iii) in the form of a ready-made micro enema; or (iv) in the form of a dry micro enema formulation to be reconstituted in a suitable solvent just prior to administration.

Liquid preparations may be prepared in the form of syrups or suspensions, e.g. solutions or
20 suspensions containing the active ingredient and the remainder consisting, for example, of sugar or sugar alcohols and a mixture of ethanol, water, glycerol, propylene glycol and polyethylene glycol. If desired, such liquid preparations may contain colouring agents, flavouring agents, preservatives, saccharine and carboxymethyl cellulose or other thickening agents. Liquid preparations may also be prepared in the form of a dry powder to
25 be reconstituted with a suitable solvent prior to use.

Solutions for parenteral administration may be prepared as a solution of a compound of the invention in a pharmaceutically acceptable solvent. These solutions may also contain stabilizing ingredients and/or buffering ingredients. Solutions for parenteral administration
30 may also be prepared as a dry preparation to be reconstituted with a suitable solvent before use.

The typical daily dose of the active substance varies within a wide range and will depend on various factors such as for example the individual requirement of each patient, the route of administration and the disease. In general, oral and parenteral dosages will be in the range of 0.1 to 1000 mg per day of active substance.

5

Medical and pharmaceutical use

The compounds of the invention are inhibitors of carboxypeptidase U either as such or, in the case of prodrugs, after administration. The compounds of the invention are thus
10 expected to be useful in those conditions where inhibition of carboxypeptidase U is beneficial, such as in the treatment or prophylaxis of thrombosis and hypercoagulability in blood and tissues of mammals, including man.

It is known that hypercoagulability may lead to thrombo-embolic diseases. Conditions
15 associated with hypercoagulability and thrombo-embolic diseases which may be mentioned include protein C resistance and inherited or acquired deficiencies in antithrombin III, protein C, protein S and heparin cofactor II. Other conditions known to be associated with hypercoagulability and thrombo-embolic disease include circulatory and septic shock, circulating antiphospholipid antibodies, homocysteinemia, heparin induced
20 thrombocytopenia and defects in fibrinolysis. The compounds of the invention are thus indicated both in the therapeutic and/or prophylactic treatment of these conditions. The compounds of the invention are further indicated in the treatment of conditions where there is an undesirable excess of proCPU/CPU.

25 Particular disease states which may be mentioned include the therapeutic and/or prophylactic treatment of venous thrombosis and pulmonary embolism, arterial thrombosis (e.g. in myocardial infarction, unstable angina, thrombosis-based stroke and peripheral arterial thrombosis) and systemic embolism usually from the atrium during arterial fibrillation or from the left ventricle after transmural myocardial infarction.

30

Moreover, the compounds of the invention are expected to have utility in prophylaxis of re-occlusion and restenosis (*i.e.* thrombosis) after thrombolysis, percutaneous trans-luminal

angioplasty (PTA) and coronary bypass operations; the prevention of re-thrombosis after microsurgery and vascular surgery in general.

Further indications include the therapeutic and/or prophylactic treatment of disseminated intravascular coagulation caused by bacteria, multiple trauma, intoxication or any other mechanism, fibrinolytic treatment when blood is in contact with foreign surfaces in the body, such as vascular grafts, vascular stents, vascular catheters, mechanical and biological prosthetic valves or any other medical device, and fibrinolytic treatment when blood is in contact with medical devices outside the body, such as during cardiovascular surgery using a heart-lung machine or in haemodialysis.

The compounds of the invention may also be combined and/or coadministered with any antithrombotic agent with a different mechanism of action, such as the antiplatelet agents acetylsalicylic acid ticlopidine, clopidogrel, thromboxane receptor and/or synthetase inhibitors, fibrinogen receptor antagonists, prostacyclin mimetics and phosphodiesterase inhibitors and ADP-receptor (P₂T) antagonists and thrombin inhibitors.

The compounds of the invention may further be combined and/or coadministered with thrombolytics such as tissue plasminogen activator (natural, recombinant or modified), streptokinase, urokinase, prourokinase, anisoylated plasminogen-streptokinase activator complex (APSAC), animal salivary gland plasminogen activators, and the like, in the treatment of thrombotic diseases, in particular myocardial infarction and stroke.

In vitro experiments

The inhibiting effect of the compounds of the present invention was estimated using the assay described in: Dirk Hendriks, Simon Scharpé and Marc van Sande, Clinical Chemistry, 31, 1936-1939 (1985); and Wei Wang, Dirk F. Hendriks, Simon S. Scharpé, The Journal of Biological Chemistry, 269, 15937-15944 (1994).

EXAMPLES

General Experimental Procedures

5 Mass spectra were recorded on a Finnigan MAT TSQ 700 triple quadrupole mass spectrometer equipped with an electrospray interface (FAB-MS) and VG Platform II mass spectrometer equipped with an electrospray interface (LC-MS). ^1H NMR and ^{13}C NMR measurements were performed on Varian UNITY plus 400, 500 and 600 spectrometers, operating at ^1H frequencies of 400, 500 and 600 MHz respectively. Chemical shifts are
10 given in ppm with the solvent as internal standard. Organic extracts were dried using MgSO_4 or Na_2SO_4 as the drying agent. Chromatography separations were performed using Merck Silica gel 60 (0.063-0.200 mm). HPLC separations were performed on a HIGHCROM KR100-10C8 column.

15 Example 15-Amino-2-[(1-benzyloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoyloxy]-pentanoic acid(a) 5-*tert*-Butoxycarbonylamino-2-hydroxy-pentanoic acid

20 Di-*t*-butyl dicarbonate (30.8 g, 0.141 mol) was added in portions during 5 min to a solution of 5-amino-2-hydroxy-pentanoic acid (17.0 g, 0.128 mol) in 0.5 M NaOH (240 mL) and dioxan (240 mL) at 5°C. The mixture was stirred for 2.5 h at room temperature. During this time 0.5 M NaOH was added to maintain pH 9-10. The dioxan was removed under reduced pressure and the aqueous phase was washed with diethyl ether. The aqueous phase was
25 acidified to pH 2-3 with KHSO_4 and extracted with ethyl acetate (3 x 300 mL). The pooled organic phases was washed with water and brine, dried and concentrated under reduced pressure to give crude 5-*tert*-butoxycarbonylamino-2-hydroxy-pentanoic acid (22.0 g, 73.7%).

30 (b) 5-*tert*-Butoxycarbonylamino-2-hydroxy-pentanoic acid methyl ester

A solution of methyl iodide (11.5 mL, 0.189 mol) in DMF (50 mL) was added dropwise during 15 min. to a mixture of 5-*tert*-butoxycarbonylamino-2-hydroxy-pentanoic acid

(22.0 g, 94.4 mmol) and NaHCO_3 (11.8 g, 141 mmol) in DMF (150 mL). After stirring over night, water was added and the mixture was extracted with ethyl acetate. The pooled organic phases were washed with water and brine, dried and concentrated under reduced pressure. The crude product was purified using chromatography (heptane/ethyl acetate, 1:1) to give 5-*tert*-butoxycarbonylamino-2-hydroxy-pentanoic acid methyl ester 9.9 g, 42.5 %)

(c) 2-[(1-Benzyloxycarbonylamino-2-methyl-propyl)-methoxy-phosphinoyloxy]-5-*tert*-butoxycarbonylamino-pentanoic acid methyl ester

A solution of PyBOP (2.1 g, 4.0 mmol) in DMF (3 mL) was added to a mixture of (1-benzyloxycarbonylamino-2-methyl-propyl)-phosphonic acid monomethyl ester (1.0 g, 3.32 mmol) and 5-*tert*-butoxycarbonylamino-2-hydroxy-pentanoic acid methyl ester (865 mg, 3.5 mmol) in DMF (4 mL) under argon. DIPEA (2.28 mL, 13.3 mmol) was added dropwise and the mixture was stirred over night. Ethyl acetate was added and the mixture was washed with 10 % KHSO_4 , satd. NaHCO_3 and brine and dried. Concentration under reduced pressure followed by chromatography (heptane/EtOAc, 1:1 \rightarrow 1:6) gave 2-[(1-benzyloxycarbonylamino-2-methyl-propyl)-methoxy-phosphinoyloxy]-5-*tert*-butoxycarbonylamino-pentanoic acid methyl ester (1.21 g, 69 %).

(d) 2-[(1-Benzyloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoyloxy]-5-*tert*-butoxycarbonylamino-pentanoic acid

1 M LiOH (5 mL) was added to a solution of 2-[(1-benzyloxycarbonylamino-2-methyl-propyl)-methoxy-phosphinoyloxy]-5-*tert*-butoxycarbonylamino-pentanoic acid methyl ester (187 mg, 0.35 mmol) in acetonitrile (5 mL). The mixture was stirred at 50°C over night and concentrated under reduced pressure. The crude product was purified using chromatography (iPrOH/conc. aq. $\text{NH}_3/\text{H}_2\text{O}$, 4:2:1) to yield 2-[(1-benzyloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoyloxy]-5-*tert*-butoxycarbonylamino-pentanoic acid (180 mg, 100 %).

(e) 5-Amino-2-[(1-benzyloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoyloxy]-pentanoic acid

TFA (3 mL) was added to a solution of 2-[(1-benzyloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoyloxy]-5-*tert*-butoxycarbonylamino-pentanoic acid (150 mg, 0.3 mmol) in methylene chloride/acetonitrile (1:1, 15 mL). The solution was stirred for 120 min and concentrated under reduced pressure to give the title compound as the TFA salt (174 mg, 100 %).

¹H NMR (500 MHz, CD₃OD): δ 1.02 (t, 6H), 1.66-2.0 (m, 4H), 2.23 (m, 1H), 2.93 (m, 2H), 3.91 (m, 1H), 4.85 (bs, 1H), 5.12 (m, 2H), 7.28-7.42 (m, 5H).

MS (+) 403.3 (M+1).

Example 2

2-[(1-Benzylloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoyloxy]-5-guanidino-pentanoic acid

(a) 2-[(1-Benzylloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoyloxy]-5-guanidino-pentanoic acid

A solution of S-methylisothiurea hydrogen sulfate (25 mg, 90 μmol) in 1 M NaOH (0.18 mL) was added to a solution of 5-amino-2-[(1-benzyloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoyloxy]-pentanoic acid (36 mg, 90 μmol) and 1 M NaOH (0.18 mL) in water/MeOH (1:1, 0.4 mL). The reaction mixture was stirred at 50°C for 6 h and concentrated under reduced pressure. The crude product was purified using HPLC (0-50 % acetonitrile, 0.1% TFA in water) to give the title compound as the TFA salt (19 mg, 38 %).

¹H NMR (500 MHz, CD₃OD): δ 1.02 (t, 6H), 1.60-1.98 (m, 4H), 2.23 (m, 1H), 3.20 (m, 2H), 3.91 (m, 1H), 4.82 (bs, 1H), 5.11 (m, 2H), 7.26-7.42 (m, 5H).

MS (+) 445 (M+1).

Example 3

5-Amino-2-[[1-(2-benzyloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy]-pentanoic acid

(a) 2-[[1-(2-Benzylloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy]-5-*tert*-butoxycarbonylamino-pentanoic acid methyl ester

Thionyl chloride (49 μ L, 0.67 mmol) was added dropwise to a solution of [1-(2-benzyloxy-carbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-phosphonic acid (208 mg, 0.48 mmol) in DMF (5 mL) at -20°C under argon. The mixture was stirred for 35 min at -5°C. A solution of 5-*tert*-Butoxycarbonylamino-2-hydroxy-pentanoic acid methyl ester
5 (166 mg, 0.67 mmol) in DMF (1 mL) was added and the mixture was stirred for 90 min at room temperature. Ethyl acetate was added and the mixture was washed with 1 M HCl, dried and concentrated under reduced pressure. The crude product was purified using chromatography (CHCl₃/MeOH/H₂O, 10:1:0 \rightarrow 10:5:1) to give 2-{[1-(2-benzyloxy-carbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy}-5-
10 *tert*-butoxycarbonylamino-pentanoic acid methyl ester (211 mg, 66 %).

(b) 2-{[1-(2-Benzylloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy}-5-*tert*-butoxycarbonylamino-pentanoic acid

1 M LiOH (3.5 mL) was added to a solution of 2-{[1-(2-benzyloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy}-5-*tert*-butoxy-carbonylamino-pentanoic acid methyl ester (211 mg, 0.32 mmol) in acetonitrile (3.5 mL)
15 and the mixture was stirred for 3 h. Ethyl acetate was added and the mixture was washed with 1 M HCl, dried and concentrated to give crude 2-{[1-(2-benzyloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy}-5-*tert*-butoxy-carbonylamino-pentanoic acid (208 mg, 100 %).
20

(c) 5-Amino-2-{[1-(2-benzyloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy}-pentanoic acid

TFA (5 mL) was added to a solution of 2-{[1-(2-benzyloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy}-5-*tert*-butoxycarbonyl-amino-pentanoic acid (208 mg, 0.32 mmol) in acetonitrile (5 mL) and the mixture was
25 stirred for 90 min. The reaction mixture was concentrated under reduced pressure to give the crude title compound as the TFA salt (212 mg, 100 %). 20 mg of the crude title compound was purified using chromatography (iPrOH/conc. aq. NH₃/H₂O, 4:2:1) to give
30 the title compound as the TFA salt (19 mg, 94 %).

¹H NMR (500 MHz, CD₃OD): δ 0.85-0.95 (m, 6H), 1.70-2.0 (m, 4H), 2.05-2.13 (m, 1H), 2.85-3.05 (m, 2H), 3.05-3.12 (m, 1H), 4.10 (bs, 1H), 4.55 (m, 1H), 4.90 (m, 1H), 5.09 (s, 2H), 7.20-7.35 (m, 10H).

5 Example 4

2-[[1-(2-Benzyloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy]-5-guanidino-pentanoic acid

(a) 5-Amino-2-hydroxy-pentanoic acid methyl ester

10 TFA (2 mL) was added to a solution of 5-*tert*-butoxycarbonylamino-2-hydroxy-pentanoic acid methyl ester in methylene chloride (10 mL) and the mixture was stirred for 3 h and then concentrated under reduced pressure to give crude 5-amino-2-hydroxy-pentanoic acid methyl ester (1 g).

15 (b) 5-(Guanidino-ω, ω'-bis(*tert*-Butoxycarbonyl)-2-hydroxy-pentanoic acid methyl ester

To a solution of 5-amino-2-hydroxy-pentanoic acid methyl ester (0.5 g, 2.0 mmol) in acetonitrile (5 mL) was added *tert*-butoxycarbonylimino-pyrazol-1-yl-methyl)-carbamic acid *tert*-butyl ester (0.77g, 2.5 mmol) followed by DIPEA (0.86 mL, 5 mmol). After stirring for 60 min ethyl acetate was added. The mixture was washed with 1 M HCl, satd.
20 NaHCO₃ and brine, dried and concentrated under reduced pressure. The crude product was purified using chromatography (heptane/ethyl acetate, 1:0 → 1:3) to give 5-(guanidino-ω, ω'-bis(*tert*-butoxycarbonyl)-2-hydroxy-pentanoic acid methyl ester (0.27 g, 35 %).

(c) 2-[[1-(2-Benzyloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy]-5-[guanidino-ω, ω'-bis(*tert*-Butoxycarbonyl)]-pentanoic acid methyl ester
25

Thionyl chloride (70 μL, 0.97 mmol) was added dropwise to a solution of [1-(2-benzyloxy-carbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-phosphonic acid (301 mg, 0.69 mmol) in DMF (5 mL) at -20°C under argon. The mixture was stirred for 20 min at
30

-5°C. A solution of 5-(guanidino- ω , ω' -bis(*tert*-Butoxycarbonyl)-2-hydroxy-pentanoic acid methyl ester (270 mg, 0.69 mmol) in DMF (1 mL) was added and the mixture was stirred for 180 min at room temperature. Ethyl acetate was added and the mixture was washed with 1 M HCl, dried and concentrated under reduced pressure. The crude product was purified using chromatography (toluene/ethyl acetate, 1:1 \rightarrow 0:1) to give 2-{{1-(2-benzyloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy}-5-[guanidino- ω , ω' -bis(*tert*-butoxycarbonyl)]-pentanoic acid methyl ester (0.27 g, 48 %).

(d) 2-{{1-(2-Benzylloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy}-5-[guanidino- ω , ω' -bis(*tert*-Butoxycarbonyl)]-pentanoic acid

A solution of LiOH (42 mg, 1.0 mmol) in water (1.0 mL) was added to a solution of 2-{{1-(2-benzyloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy}-5-[guanidino- ω , ω' -bis(*tert*-butoxycarbonyl)]-pentanoic acid methyl ester (160 mg, 0.2 mmol) in acetonitrile (1.0 mL) and the mixture was stirred for 15 min. Ethyl acetate was added and the mixture was washed with 1 M HCl and brine, dried and concentrated to give crude 2-{{1-(2-benzyloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy}-5-[guanidino- ω , ω' -bis(*tert*-butoxycarbonyl)]-pentanoic acid (160 mg, 100 %).

(e) 2-{{1-(2-Benzylloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy}-5-guanidino-pentanoic acid

TFA (2 mL) was added to a solution of 2-{{1-(2-benzyloxycarbonylamino-3-phenyl-propionylamino)-2-methyl-propyl]-hydroxy-phosphinoyloxy}-5-[guanidino- ω , ω' -bis(*tert*-butoxycarbonyl)]-pentanoic acid (160 mg, 0.2 mmol) in acetonitrile (5 mL) and the mixture was stirred for 60 min and then concentrated under reduced pressure. The crude product was purified using chromatography (iPrOH/conc. aq. NH₃/H₂O, 4:2:1) to give the title compound as the TFA salt (30 mg, 21 %).

¹H NMR (500 MHz, CD₃OD): δ 0.80-0.98 (m, 6H), 1.53-1.95 (m, 4H), 2.01-2.30 (m, 1H), 2.90 (m, 1H), 3.10-3.30 (m, 3H), 3.94-4.10 (m, 1H), 4.41-4.55 (m, 1H), 4.68 (m, 1H), 5.03 (m, 2H), 7.18-7.37 (m, 5H).

MS (+) 592 (M+1).

Example 5

2-Hydroxycarbamoyl-4-piperidin-4-yl-butvric acid

5

(a) Piperidin-4-yl-acetic acid

Pyridin-4-yl-acetic acid hydrochloride (20.0 g, 115 mmol) was added to water/25 % ammonia (125 mL:10mL). The mixture was degassed and flushed with nitrogen before addition of rhodium on activated alumina (0.45 g). The mixture was again degassed, then
10 stirred in a hydrogen atmosphere at 50 bar for 16 h. Filtration of the reaction mixture through filter paper afforded the bulk of the catalyst which was recycled after washing with methanol. The filtrate was then filtered through Celite and concentrated to afford a white solid (19.7 g, 96 % yield).

15 (b) 4-Carboxymethyl-piperidine-1-carboxylic acid *tert*-butyl ester

To a solution of piperidin-4-yl-acetic acid (19.7 g, 148 mmol) in THF-water (417 mL, 1:1) was added di-*tert*-butyl dicarbonate (32.3 g, 148 mmol) and sodium bicarbonate (12.5 g, 148 mmol), and the reaction stirred at room temperature for 16 h. THF was then removed under reduced pressure and the aqueous phase extracted with dichloromethane and the
20 organic layer discarded. The aqueous layer was then acidified to pH 1-2 with 1 M HCl solution and extracted with ethyl acetate. The organic phase was washed with brine, dried and concentrated under reduced pressure to give 4-carboxymethyl-piperidine-1-carboxylic acid *tert*-butyl ester (16.7 g, 46 %).

25 (c) 4-(2-Hydroxy-ethyl)-piperidine-1-carboxylic acid *tert*-butyl ester

To a solution of 4-carboxymethyl-piperidine-1-carboxylic acid *tert*-butyl ester (16.7 g, 69.0 mmol) in THF (100 mL) was added diborane (151 mL, 1.0 M solution in THF) over a period of 10 min at 0°C. Hydrogen gas was rapidly evolved and after gas evolution had ceased the reaction was stirred at room temperature for 1 h. The reaction mixture was again
30 cooled to 0°C, and 1 M aqueous HCl was added dropwise to the reaction mixture with further evolution of hydrogen. Addition of HCl was continued until the evolution of hydrogen had almost ceased. The mixture was then stirred for 10 min and made basic (pH

13-14) by the addition of 1 M NaOH solution. The aqueous solution was extracted with ethyl acetate, the organic phase washed with brine, dried and concentrated under reduced pressure to yield 4-(2-hydroxy-ethyl)-piperidine-1-carboxylic acid *tert*-butyl ester (15.2 g, 97 %).

5 (d) 4-(2-Oxo-ethyl)-piperidine-1-carboxylic acid *tert*-butyl ester

Periodinane (36.1 g, 85.2 mmol) was added to 4-(2-hydroxy-ethyl)-piperidine-1-carboxylic acid *tert*-butyl ester (15.0 g, 65.5 mmol) in CH₂Cl₂ (230 mL) and stirred for 90 min.

Diethyl ether (560 ml) was added and precipitates were removed by extraction with 10 %

10 Na₂S₂O₃/saturated NaHCO₃ (1:1, 350 mL). The organic layer was washed with 0.5 M NaOH solution and brine. The organic phase was dried and concentrated under reduced pressure to yield 4-(2-Oxo-ethyl)-piperidine-1-carboxylic acid *tert*-butyl ester (8.50 g, 57 %).

15 (e) 4-[2-(2,2-Dimethyl-4,6-dioxo-[1,3]dioxan-5-yl)-ethyl]-piperidine-1-carboxylic acid *tert*-butyl ester.

To a solution of meldrums acid (1.68 g, 11.66 mmol) and 4-(2-oxo-ethyl)-piperidine-1-carboxylic acid *tert*-butyl ester (2.21 g, 9.72 mmol) in dichloromethane (40 mL) was added acetic acid (0.055 mL, 0.972 mmol) and piperidine (0.096 mL, 0.972 mmol). The mixture
20 was heated at reflux for 3 h, and then allowed to attain room temperature. After being diluted with *tert*-butyl methyl ether, the mixture was washed with NaHCO₃ (sat.) and brine. The organic phase was dried, filtered and concentrated. The residue was dissolved in a mixture of EtOH (40 mL) and acetic acid (20 mL). The solution was cooled to 0°C, and NaBH₄ (0.554 g, 14.6 mmol) was added in portions after which the solution was allowed to
25 stir for 30 min at rt and then acidified to pH 3 with 1 M HCl. The solution was extracted several times with dichloromethane. The combined organic phases were dried, filtered, concentrated and filtered through a pad of silica gel (dichloromethane). The solvent was evaporated to give 4-[2-(2,2-Dimethyl-4,6-dioxo-[1,3]dioxan-5-yl)-ethyl]-piperidine-1-carboxylic acid *tert*-butyl ester as a colorless oil (2.30 g, 65 %) which solidified on
30 standing.

(f) 2-Hydroxycarbamoyl-4-piperidin-4-yl-butvic acid.

A GC-autosample vial (2 mL) equipped with a septum cap and a small stirbar was charged with 4-[2-(2,2-Dimethyl-4,6-dioxo-[1,3]dioxan-5-yl)-ethyl]-piperidine-1-carboxylic acid *tert*-butyl ester (17.8 mg, 0.05 mmol) and flushed with nitrogen. N.O-Bis(trimethylsilyl)-hydroxylamine (0.2 mL) was added via syringe and the resulting solution was stirred at rt over night. The mixture was concentrated under vacuum and the residue dissolved in dichloromethane/MeOH (4:1) and applied onto a small plug of ion exchange resin (200 mg, isolute™, aminoresin), washed with dichloromethane/MeOH (4:1) and then eluted with dichloromethane/MeOH/AcOH (3:1:1). The eluate was concentrated, the residue dissolved in dichloromethane/TFA (1:1, 2 mL) and stirred for 1h at room temperature. Evaporation of the solvent gave the title compound as the TFA salt (16 mg, 93 %) as a colourless oil.

¹H NMR (600 MHz, CD₃OD) δ 1.21-1.40 (m, 4H), 1.53-1.62 (m, 1H), 1.80-1.99 (m, 4H), 2.90-2.98 (m, 2H), 3.06 (t, 1H), 3.32-3.39 (m, 2H).

M (+) 231 (M+1).

Example 6

N-hydroxy-2-piperidin-3-ylmethyl-malonamic acid

(a) 3-Hydroxymethyl-piperidine-1-carboxylic acid *tert*-butyl ester

3-Hydroxymethyl-piperidine (20.0 g, 0.17 mmol) in acetonitrile was treated with di-*tert*-butyl dicarbonate (37.9 g, 0.17 mol) and DMAP (2.13 g, 1.74 mmol). The reaction mixture was stirred at ambient temperature for 5 h and then concentrated under reduced pressure. The crude product was purified by flash chromatography (hexane/EtOAc, 70:30) to give 3-hydroxymethyl-piperidine-1-carboxylic acid *tert*-butyl ester (16.0 g, 44 %).

(b) 3-Formyl-piperidine-1-carboxylic acid *tert*-butyl ester

Periodinane (18.2 g, 42.9 mmol) was added to 3-hydroxymethyl-piperidine-1-carboxylic acid *tert*-butyl ester (7.10 g, 33.0 mmol) in CH₂Cl₂ (230 mL) and stirred for 90 min. Diethyl ether (230 mL) was added and precipitates were removed by extraction with 10 % Na₂S₂O₃/saturated NaHCO₃ (1:1, 230 mL). The organic layer was washed with 0.5 M NaOH solution and brine. The organic phase was dried and concentrated under reduced pressure to yield 3-Formyl-piperidine-1-carboxylic acid *tert*-butyl ester (6.50 g, 93 %).

(c) N-Hydroxy-2-piperidin-3-ylmethyl-malonamic acid

The title compound was prepared from 3-formyl-piperidine-1-carboxylic acid *tert*-butyl ester by the method described in Example 5. Yield: (50 %).

5 ¹H NMR (600 MHz, CD₃OD) δ 1.18-1.30 (m, 1H), 1.61-1.99 (m, 6H), 2.64 (t, 1H), 2.86 (t, 1H), 3.20-3.38 (m, 3H).

M (+) 217 (M+1).

Example 7

10 2-(6-Amino-pyridin-3-ylmethyl)-N-hydroxy-malonamic acid.

The title compound was prepared from (5-Formyl-pyridin-2-yl)-carbamic acid *tert*-butyl ester by the method described in Example 5. Yield: (23 %).

¹H NMR (600 MHz, CD₃OD) δ 2.99-3.10 (m, 2H), 3.36-4.01 (m, 1H), 6.94 (d, 1H), 7.64 (s, 1H), 7.82 (d, 1H).

15 M (+) 226 (M+1).

Example 8

2-(2-Amino-pyridin-4-ylmethyl)-N-hydroxy-malonamic acid.

20 (a) (4-Formyl-pyridin-2-yl)-carbamic acid tert-butyl ester

(4-Hydroxymethyl-pyridin-2-yl)-carbamic acid *tert*-butyl ester (1.91 g, 8.51 mmol) was dissolved in dry DMSO (10 mL) and the reaction flask immersed in a waterbath at 15°C. Triethylamine (1.72 g, 17.0 mmol) was added followed by sulfur trioxide pyridine complex (2.41 g, 15.1 mmol). The reaction mixture was stirred for 2 h and poured onto
25 crushed ice and the product extracted with CHCl₃. The organic extract was washed with water, dried concentrated under reduced pressure. The crude product was purified by flash chromatography (hexane/EtOAc, 80:20) to give (4-Formyl-pyridin-2-yl)-carbamic acid *tert*-butyl ester (1.57 g, 83 %).

30 (b) 2-(2-Amino-pyridin-4-ylmethyl)-N-hydroxy-malonamic acid

The title compound was prepared from (4-Formyl-pyridin-2-yl)-carbamic acid *tert*-butyl ester by the method described in Example 5. Yield: (48 %).

¹H NMR (600 MHz, CD₃OD) δ 3.10 (dd, 1H), 3.19 (dd, 1H), 3.47 (dd, 1H), 6.77 (d, 1H), 7.82 (s, 1H), 7.71 (d, 1H).

M (+) 226 (M+1).

5 Example 9

2-[2-(1-*tert*-Butoxycarbonyl-piperidin-4-yl)-ethyl]-malonic acid.

To a solution of 4-[2-(2,2-Dimethyl-4,6-dioxo-[1,3]dioxan-5-yl)-ethyl]-piperidine-1-carboxylic acid *tert*-butyl ester (17.8 mg, 0.05 mmol) in acetic acid (1 mL) was added 6 M HCl (2 mL). The solution was stirred at room temperature over night and then concentrated
10 to yield the title compound as the hydrochloric acid salt (15 mg, 100 %).

¹H NMR (600 MHz, CD₃OD) δ 1.30-1.40 (m, 4H), 1.57-1.64 (m, 1H), 1.83-1.90 (m, 1H), 1.90-1.98 (m, 1H), 3.31-3.39 (m, 2H).

M (+) 216 (M+1).

15 Example 10

2-[2-(1-*tert*-Butoxycarbonyl-piperidin-3-yl)-methyl]-malonic acid.

The title compound was prepared from 3-(2,2-Dimethyl-4,6-dioxo-[1,3]dioxan-5-ylmethyl)-piperidine-1-carboxylic acid *tert*-butyl ester by the method described in Example 9. Yield: (100 %).

20 ¹H NMR (600 MHz, CD₃OD) δ 1.20-1.30 (m, 1H), 1.63-1.76 (m, 1H), 1.78-1.97 (m, 5H), 2.65 (t, 1H), 2.83-2.92 (m, 1H), 3.29-3.38 (m, 2H), 3.42-3.48 (m, 1H).

M (+) 202 (M+1).

Example 11

25 2-[2-(1-*tert*-Butoxycarbonyl-piperidin-4-yl)-methyl]-malonic acid

(a) Piperidine-1,4-dicarboxylic acid mono-*tert*-butyl ester

Piperidine-4-carboxylic acid (24.5 g, 0.19 mmol) in THF/water (1:1, 417 mL) was treated with di-*tert*-butyl dicarbonate (41.49 g, 0.19 mol) and sodium bicarbonate (16.0 g, 0.19
30 mol). The reaction mixture was stirred at ambient temperature for 16 h. The THF was then removed under reduced pressure and the aqueous phase washed with dichloromethane. The aqueous layer was then acidified to pH 1-2 with 1 M HCl solution and extracted with ethyl

acetate. The organic phase was washed with brine and dried to give piperidine-1,4-dicarboxylic acid mono-*tert*-butyl ester (35.9 g, 83 %).

(b) 4-Hydroxymethyl-piperidine-1-carboxylic acid *tert*-butyl ester

5 To a solution of piperidine-1,4-dicarboxylic acid mono-*tert*-butyl ester (19.3 g, 84.0 mmol) in THF (100 mL) was added diborane (185 mL, 1.0 M solution in THF) over a period of 10 min at 0°C. Hydrogen gas was rapidly evolved and after gas evolution had ceased the reaction was stirred at room temperature for 1 h. The reaction mixture was again cooled to 0°C, and 1 M aqueous HCl was added dropwise to the reaction mixture with further evolution of
10 hydrogen. Addition of HCl was continued until the evolution of hydrogen had almost ceased. The mixture was then stirred for 10 min and made basic (pH 13-14) by the addition of 1 M NaOH solution. The aqueous solution was extracted with ethyl acetate, the organic phase washed with brine, dried and concentrated under reduced pressure to yield 4-Hydroxymethyl-piperidine-1-carboxylic acid *tert*-butyl ester (18.12 g, 100 %).

15

(c) 4-Formyl-piperidine-1-carboxylic acid *tert*-butyl ester

Periodinane (26.9 g, 63.5 mmol) was added to 4-Hydroxymethyl-piperidine-1-carboxylic acid *tert*-butyl ester (10.5 g, 48.8 mmol) in CH₂Cl₂ (200 mL) and stirred for 90 min. Diethyl ether (560 mL) was added and precipitates were removed by extraction with 10 %
20 Na₂S₂O₃/saturated NaHCO₃ (1:1, 300 mL). The organic layer was washed with 0.5 M NaOH solution and brine. The organic phase was dried and concentrated under reduced pressure. Purification using flash chromatography (hexane/EtOAc, 8:2) gave 4-Formyl-piperidine-1-carboxylic acid *tert*-butyl ester (8.5 g, 81 %).

25 (d) 2-[2-(1-*tert*-Butoxycarbonyl-piperidin-4-yl)-methyl]-malonic acid.

The title compound was prepared from 4-Formyl-piperidine-1-carboxylic acid *tert*-butyl ester by the method described in Example 5 and 9. Yield: (100 %).

¹H NMR (600 MHz, CD₃OD) δ 1.38-1.48 (m, 2H), 1.61-1.75 (m, 1H), 1.82-1.90 (m, 2H), 1.92-2.02 (m, 2H), 2.90-3.01 (m, 2H), 3.35-3.42 (m, 2H), 3.42-3.48 (m, 1H).

30 M (+) 202 (M+1).

Example 12

2-(6-*tert*-Butoxycarbonylamino-pyridin-3-ylmethyl)-malonic acid.

The title compound was prepared from [5-(2,2-Dimethyl-4,6-dioxo-[1,3]dioxan-5-ylmethyl)-pyridin-2-yl]-carbamic acid *tert*-butyl ester by the method described in Example 9. Yield: (100 %).

5 ¹H NMR (600 MHz, CD₃OD) δ 3.08 (d, 2H), 3.66 (t, 1H), 6.98 (d, 1H), 7.73 (s, 1H), 7.92 (d, 1H).

M (+) 211 (M+1).

Example 13

10 2-(2-*tert*-Butoxycarbonylamino-pyridin-4-ylmethyl)-malonic acid.

The title compound was prepared from [4-(2,2-Dimethyl-4,6-dioxo-[1,3]dioxan-5-ylmethyl)-pyridin-2-yl]-carbamic acid *tert*-butyl ester by the method described in Example 9. Yield: (100 %).

¹H NMR (600 MHz, CD₃OD) δ 3.10 (d, 2H), 3.79 (t, 1H), 6.84 (d, 1H), 7.92 (s, 1H), 7.77 (d, 1H).

15 M (+) 211 (M+1).

Example 142-(2-Amino-pyridin-4-ylmethyl)-succinic acid

20

(a) 2-(2-*tert*-Butoxycarbonylamino-pyridin-4-ylmethylene)-succinic acid 4-benzyl ester 1-*tert*-butyl ester

Butyllithium (1.6 M in hexane, 14.8 ml, 23.7 mmol) was added dropwise to a solution of 2-(diethoxy-phosphoryl)-succinic acid 4-benzyl ester 1-*tert*-butyl ester (9.50 g, 23.7 mmol) in THF (75 mL) at 0°C under nitrogen. After stirring at 0°C for 1 h, the solution was transferred to a solution of (4-Formyl-pyridin-2-yl)-carbamic acid *tert*-butyl ester (3.70 g, 16.6 mmol) in THF (75 mL). The resulting reaction mixture was stirred at 0°C for 1 h before being allowed to warm to 25 °C, and the mixture was stirred overnight. Water (400 mL) was added and the product extracted with CH₂Cl₂ (3x50 mL). The combined organic layers were dried and concentrated. Flash chromatography (hexane/EtOAc, 4:1) gave 2-(2-*tert*-Butoxycarbonylamino-pyridin-4-ylmethylene)-succinic acid 4-benzyl ester 1-*tert*-butyl ester 4.30 g (55 %).

25
30

(b) 2-(2-*tert*-Butoxycarbonylamino-pyridin-4-ylmethyl)-succinic acid 1-*tert*-butyl ester
2-(2-*tert*-Butoxycarbonylamino-pyridin-4-ylmethylene)-succinic acid 4-benzyl ester 1-*tert*-
butyl ester (2.81 g, 7.60 mmol) and Pd/C (10 %, 400 mg) were suspended in EtOH and
5 hydrogenated at 41 atm. and 28°C for 3 days. The catalyst was removed from the reaction
mixture by filtration. The catalyst was washed with EtOH (96 %). 1 M K₂CO₃ (30 mL) was
added to the filtrate followed by addition of water (50 mL). After 2 days the reaction
mixture was evaporated to ca 80 mL, then brine (10 mL) was added and the reaction
mixture extracted with ether. The aqueous phase was acidified to pH=3 and extracted with
10 chloroform. Methanol (25 mL) was added and the reaction mixture was dried (Na₂SO₄ +
CaSO₄) and filtered to give 2-(2-*tert*-butoxycarbonylamino-pyridin-4-ylmethyl)-succinic
acid 1-*tert*-butyl ester (1.90 g, 83 %).

(c) 2-(2-Amino-pyridin-4-ylmethyl)-succinic acid
15 To a solution of 2-(2-*tert*-butoxycarbonylamino-pyridin-4-ylmethyl)-succinic acid 1-*tert*-
butyl ester (164 mg, 0.43 mmol) in methylene chloride (1.5 mL) was added TFA (1.5 mL).
The reaction mixture was stirred for 2.5 h and then concentrated under reduced pressure.
The residue was lyophilized to give the title compound as the TFA salt (126mg, 87 %)
¹H NMR (500 MHz, D₂O): δ 2.58-2.80 (m, 2H), 2.88-3.07 (m, 2H), 3.13-3.26 (m, 1H),
20 6.79-6.84 (dd, 1H), 6.84-6.88 (s, 1H), 7.69-7.75 (d, 1H).
MS (+) 225 (M+1).

Example 15

trans-2-(4-Amino-cyclohexylmethyl)-succinic acid

25

(a) 4-[*N*-(*tert*-Butoxycarbonyl)amino]-cyclohexane carboxylic acid

To a solution of *cis*-4-aminocyclohexane carboxylic acid (9.90 g, 69.0 mmol) in water (120
mL) and dioxane (120 mL) was added KOH (3.73 g, 56 mmol) followed by di-*tert*-butyl
dicarbonate (15.3 g, 70.0 mmol). The reaction mixture was stirred at room temperature
overnight. Water was added and the product was extracted with CHCl₃. The combined
30 organic extracts were washed with water, dried and concentrated under reduced pressure to
give 4-[*N*-(*tert*-butoxycarbonyl)amino]-cyclohexane carboxylic acid (14.1 g, 84 %).

(b) [4-(Methoxy-methyl-carbamoyl)-cyclohexyl]-carbamic acid *tert*-butyl ester

A solution of 4-[*N*-(*tert*-butoxycarbonyl)amino]-cyclohexane carboxylic acid (11.95 g, 49.0 mmol), *O,N*-dimethylhydroxylamine (4.88 g, 50.0 mmol), DCC (9.60 g, 50 mmol) and triethylamine (5.06 g, 50.0 mmol) in DMF (150 mL) was stirred at room temperature overnight. Water (500 mL) was added and the mixture was extracted with CHCl₃. The organic phase was washed with water, dried and concentrated under reduced pressure. Purification by flash chromatography (hexane/EtOAc 1:1) gave [4-(methoxy-methyl-carbamoyl)-cyclohexyl]-carbamic acid *tert*-butyl ester (8.50 g, 61 %).

(c) (4-Formyl-cyclohexyl)-carbamic acid *tert*-butyl ester

[4-(Methoxy-methyl-carbamoyl)-cyclohexyl]-carbamic acid *tert*-butyl ester (7.50 g, 26.2 mmol) in dry ether (150 ml) was reduced with an excess LiAlH₄. The reaction mixture was quenched by careful addition of water and extracted with CHCl₃. The mixture was dried and concentrated under reduced pressure to give (4-formyl-cyclohexyl)-carbamic acid *tert*-butyl ester (6.30 g, 93 %).

(d) trans-[4-(Benzylimino-methyl)-cyclohexyl]-carbamic acid *tert*-butyl ester

A mixture of (4-Formyl-cyclohexyl)-carbamic acid *tert*-butyl ester (3.80 g, 16.7 mmol), benzylamine (1.82 g, 16.7 mmol), acetic acid (0.01 g, 16.7 mmol) and anhydrous magnesium sulfate (4.01 g, 33.3 mmol) in methylene chloride (20 mL) was stirred for 5 days. The mixture was filtered through Celite and concentrated under reduced pressure to give trans-[4-(benzylimino-methyl)-cyclohexyl]-carbamic acid *tert*-butyl ester (5.10 g, 97 %) as a 97:3 *trans*:*cis* mixture.

(e) trans-(4-Formyl-cyclohexyl)-carbamic acid *tert*-butyl ester

A solution of trans-[4-(benzylimino-methyl)-cyclohexyl]-carbamic acid *tert*-butyl ester (2.50 g, 8.00 mmol) and oxalic acid (0.80 g) in water/THF (50 mL, 1:1) was stirred for 10 h at room temperature. The reaction mixture was concentrated under reduced pressure and methylene chloride (50 mL) was added to the residue. The organic phase was dried and concentrated under reduced pressure to give trans-(4-Formyl-cyclohexyl)-carbamic acid *tert*-butyl ester (1.3 g, 80 %).

(f) trans 2-(4-*tert*-Butoxycarbonylamino-cyclohexylmethylene)-succinic acid 4-benzyl ester 1-*tert*-butyl ester

Butyllithium (1.6 M in hexane, 5.0 ml, 8.00 mmol) was added dropwise to a solution of 2-
5 (diethoxy-phosphoryl)-succinic acid 4-benzyl ester 1-*tert*-butyl ester (3.21 g, 8.00 mmol)
in THF (25 mL) at 0°C under nitrogen. After stirring at 0°C for 1 h, the solution was
transferred to a solution of trans-(4-Formyl-cyclohexyl)-carbamic acid *tert*-butyl ester
(1.30 g, 5.72 mmol) in THF (10 mL). The resulting mixture was stirred at 0°C for 1 h and
at room temperature overnight. Water was added and the product extracted with CH₂Cl₂.
10 The organic phase was dried and concentrated under reduced pressure. Flash chromatography
(hexane/EtOAc, 80:20) gave trans-2-(4-*tert*-Butoxycarbonylamino-cyclohexylmethylene)-succinic acid 4-benzyl ester 1-*tert*-butyl ester (1.10 g)

(g) trans-2-(4-*tert*-Butoxycarbonylamino-cyclohexylmethyl)-succinic acid 1-*tert*-butyl ester

A solution of trans-2-(4-*tert*-Butoxycarbonylamino-cyclohexylmethylene)-succinic acid 4-benzyl ester 1-*tert*-butyl ester (243 mg, 0.51 mmol) and palladium (5 % on charcoal) in ethanol (15 mL) was hydrogenated at 4 bar for 3 h. The catalyst was removed from the reaction mixture by filtration. The catalyst was washed with ethanol and the solution was
20 concentrated under reduced pressure to give crude trans-2-(4-*tert*-Butoxycarbonylamino-cyclohexylmethyl)-succinic acid 1-*tert*-butyl ester (217 mg, >100 %).

(h) trans-2-(4-Amino-cyclohexylmethyl)-succinic acid

To a solution of trans-2-(4-*tert*-Butoxycarbonylamino-cyclohexylmethyl)-succinic acid 1-
25 *tert*-butyl ester (200 mg, 0.52 mmol) in methylene chloride (1.32 g, 15.6 mmol) was added triethylsilane (150 mg, 1.30 mmol) followed by TFA (770 mg, 6.75 mmol). The reaction mixture was stirred for 2.5 h and then concentrated under reduced pressure. Purification by HPLC (0-80 % acetonitrile, 0.1 % TFA in water) gave the title compound as the TFA salt (60 mg, 34 %)

30 ¹H NMR (500 MHz, D₂O): δ 0.99-1.11 (m, 2H), 1.30-1.46 (m, 4H), 1.54-1.62 (m, 1H), 1.79-1.86 (m, 1H), 1.89-1.96 (m, 1H), 1.99-2.06 (m, 2H), 2.58-2.71 (m, 2H), 2.85-2.95 (m, 1H), 3.08-3.17 (m, 1H).

MS (+) 230 (M+1).

Example 16

2-(6-Amino-pyridin-3-ylmethyl)-N-benzyl-N-hydroxy-succinamic acid

5

(a) N-Benzyl-N-benzyloxy-2-(6-tert-butoxycarbonylamino-pyridin-3-ylmethyl)-succinamic acid tert-butyl ester

To a solution of 2-(2-tert-butoxycarbonylamino-pyridin-4-ylmethyl)-succinic acid 1-tert-butyl ester (0.67 g, 1.76 mmol) in CH₂Cl₂ (25 mL) was added N-benzyl-N-benzyloxy amine (0.42 g, 1.94 mmol), DCC (0.40 g, 1.94 mmol) and DMAP (0.02 g, 0.17 mmol) and the mixture was stirred overnight. Water was added and the mixture was extracted with CH₂Cl₂. The organic phase was dried and filtered and the residue purified by flash chromatography (hexane/EtOAc, 4:1) to give N-Benzyl-N-benzyloxy-2-(6-tert-butoxycarbonylamino-pyridin-3-ylmethyl)-succinamic acid tert-butyl ester (0.51 g, 50 %).

15

(b) 2-(6-Amino-pyridin-3-ylmethyl)-N-benzyl-N-benzyloxy-succinamic acid

To a solution of N-Benzyl-N-benzyloxy-2-(6-tert-butoxycarbonylamino-pyridin-3-ylmethyl)-succinamic acid tert-butyl ester (1.0 g, 1.7 mmol) in methylene chloride (10 mL) was added TFA (4 mL) at 0°C. The reaction mixture was stirred for 4 hs and then concentrated under reduced pressure to give crude 2-(6-Amino-pyridin-3-ylmethyl)-N-benzyl-N-benzyloxy-succinamic acid as the TFA salt (0.9 g, 100 %).

20

(c) 2-(6-Amino-pyridin-3-ylmethyl)-N-benzyl-N-hydroxy-succinamic acid

A solution of 2-(6-Amino-pyridin-3-ylmethyl)-N-benzyl-N-benzyloxy-succinamic acid (0.9 g, 1.7 mmol) and palladium (0.5g, 5 % on charcoal) in methanol (100 mL) was hydrogenated at 1 bar for 2 h. The reaction mixture was filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (CHCl₃/MeOH/H₂O, 10:5:1) to give the title compound (123 mg, 22 %).

25

¹H NMR (600 MHz, CD₃OD): δ 2.50-3.03 (m, 5H), 4.72 (q, 2H), 6.65 (d, 1H), 7.18-7.31 (m, 6H), 7.53 (d, 1H), 7.65 (s, 1H).

30

MS (+) 330 (M+1).

Example 172-(6-amino-pyridin-3-ylmethyl)-3-[hydroxy-(3-phenyl-propyl)-phosphinoyl]-propionic acid5 (a) 2-(6-*tert*-Butoxycarbonylamino-pyridin-3-ylmethyl)-3-[hydroxy-(3-phenyl-propyl)-phosphinoyl]-propionic acid ethyl ester

A solution of (3-Phenyl-propyl)-phosphinic acid (0.579g 3.143mmol) and 2-(6-*tert*-butoxycarbonylamino-pyridin-3-ylmethyl)-acrylic acid ethyl ester (0.327g 1.067 mmol) in MeCN (7.5 mL) was degassed using the freeze-thaw technique. BSA (2.55g
10 12.57 mmol) was added and the mixture was stirred for 3 days and concentrated under reduced pressure. The residue was dissolved in chloroform, washed with NaHCO₃ and brine, dried and concentrated under reduced pressure. Flash chromatography (CH₂Cl₂/MeOH, 8:2) gave 2-(6-*tert*-butoxycarbonylamino-pyridin-3-ylmethyl)-3-[hydroxy-(3-phenyl-propyl)-phosphinoyl]-propionic acid ethyl ester (0.240g, 16%).

15

(b) 2-(6-*tert*-Butoxycarbonylamino-pyridin-3-ylmethyl)-3-[hydroxy-(3-phenyl-propyl)-phosphinoyl]-propionic acid

To a solution of 2-(6-*tert*-butoxycarbonylamino-pyridin-3-ylmethyl)-3-[hydroxy-(3-phenyl-propyl)-phosphinoyl]-propionic acid ethyl ester (0.240g 0.489 mmol) in MeCN (3
20 mL) was added a solution of LiOH (0.059 g 2.45 mmol) in H₂O (3 mL). The mixture was then stirred at 20°C for 1.5 hours. Ethylacetate was added and the mixture was washed with 1 M HCL and brine and filtered to give 2-(6-*tert*-butoxycarbonylamino-pyridin-3-ylmethyl)-3-[hydroxy-(3-phenyl-propyl)-phosphinoyl]-propionic acid (0.174 g, 77%) as white crystals.

25

(c) 2-(6-Amino-pyridin-3-ylmethyl)-3-[hydroxy-(3-phenyl-propyl)-phosphinoyl]-propionic acid

To a mixture of 2-(6-*tert*-butoxycarbonylamino-pyridin-3-ylmethyl)-3-[hydroxy-(3-phenyl-propyl)-phosphinoyl]-propionic acid (0.170 g, 0.367 mmol) in ethylacetate (3 mL)
30 at 4°C was slowly added ethylacetat (4 mL, saturated with HCl(g)). The mixture was then stirred for 22 h. Concentration under reduced pressure gave the title compound (0.124 g, 93%) as the hydrochloride salt.

¹H NMR (300 MHz, CD₃SOCD₃): δ 1.52-1.72 (m, 4H), 1.88-1.96 (m, 2H), 2.47-2.49 (t, 1H), 2.57-2.62 (m, 2H), 2.77-2.82 (m, 2H), 6.94-6.97 (d, 1H), 7.15-7.29 (m, 4H), 7.75-7.80 (m, 2H), 8.05 (s, 1H)
MS (+) 363 (M+1).

5

Example 18

2-(6-Amino-5-methyl-pyridin-3-ylmethyl)-3-[(1-benzyloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoyl]-propionic acid

10 (a) 2-[(1-Benzyloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoylmethyl]-3-(6-bis(tert-butoxycarbonyl)amino-5-methyl-pyridin-3-yl)-propionic acid ethyl ester

A solution of 2-(6-bis(tert-butoxycarbonyl)amino-5-methyl-pyridin-3-ylmethyl)-acrylic acid ethyl ester (0.6 g, 1.43 mmol) and (1-benzyloxycarbonylamino-2-methyl-propyl)-phosphinic acid (0.678 g, 2.5 mmol) in dry acetonitrile (10 mL) was degassed using the freeze-thaw technique. Bistrimethylsilylacetamide (5 mL, 20.3 mmol) was added under argon. The resulting mixture was stirred at room temperature for 84 h and then concentrated under reduced pressure. The remaining mixture was dissolved in chloroform and washed with saturated aqueous NaHCO₃. The aqueous phase was extracted with chloroform and EtOAc. The combined organic extracts were dried and concentrated under reduced pressure. The crude product was purified by flash chromatography (EtOAc/EtOH, 100:20 → 100:25) to yield 2-[(1-benzyloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoylmethyl]-3-(6-bis(tert-butoxycarbonyl)amino-5-methyl-pyridin-3-yl)-propionic acid ethyl ester (650 mg, 65,9%)

25 (b) 2-[(1-Benzyloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoylmethyl]-3-(6-tert-butoxycarbonylamino-5-methyl-pyridin-3-yl)-propionic acid

1 M LiOH (2 mL) was added dropwise to a solution of 2-[(1-benzyloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoylmethyl]-3-(6-bis(tert-butoxycarbonyl)amino-5-methyl-pyridin-3-yl)-propionic acid ethyl ester (100 mg, 0.145 mmol) in acetonitrile (2 mL). The mixture was stirred overnight and concentrated under reduced pressure. The mixture was purified by column chromatography (isopropanol/concentrated aqueous NH₃/water, 4:2:1) to give unpure product. The unpure product was stirred with MeOH,

30

filtered and concentrated under reduced pressure. The crude product was stirred with EtOH, filtered and concentrated under reduced pressure. The crude product was stirred with ethylacetate/ethanol, filtered and concentrated under reduced pressure to give 2-[(1-benzyloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoylmethyl]-3-(6-tert-butoxycarbonylamino-5-methyl-pyridin-3-yl)-propionic acid (52 mg, 63.8 %).

(c) 2-(6-Amino-5-methyl-pyridin-3-ylmethyl)-3-[(1-benzyloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoyl]-propionic acid

A solution of 2-[(1-benzyloxycarbonylamino-2-methyl-propyl)-hydroxy-phosphinoylmethyl]-3-(6-tert-butoxycarbonylamino-5-methyl-pyridin-3-yl)-propionic acid (52 mg, 0.09 mmol) and ethylacetate (6 mL, saturated with HCl (g)) was stirred for 20 min. The mixture was concentrated under reduced pressure, washed with acetonitrile and dissolved in ethanol and petroleum ether. The solution was concentrated under reduced pressure to give the title compound (41 mg, 91.3 %) as the hydrochloride salt.

¹H NMR (600 MHz, D₂O): δ 0.99-1.05 (m, 6H), 1.75-1.85 (m, 1H), 2.10-2.28 (m, 5H), 2.68-3.15 (m, 3H), 3.75-3.81 (m, 1H), 5.04-5.19 (m, 2H), 7.20-7.37 (m, 5H), 7.51-7.71 (m, 2H).

MS (+) 501 (M+1)

Example 19

2-(6-Amino-pyridin-3-ylmethyl)-3-methyl-succinic acid

(a) 2-Ethoxycarbonyl-3-methyl-succinic acid diethyl ester

A solution of diethylmalonate (2.44g, 15 mmol) and cesium fluoride (2.32g, 15mmol) in DMF (20 mL) was stirred for 1 h at room temperature under argon. A solution of 2-methanesulfonyloxy-propionic acid ethyl ester (1.0 g, 5 mmol) in DMF (5 mL) was added and the mixture was stirred at 45°C overnight. The mixture was poured into H₂O and extracted with ethyl acetate. The organic layers were washed with water, dried, and concentrated under reduced pressure. The crude product was purified by vacuum distillation. The product was purified by column chromatography (petroleum ether/ethyl acetate, 4:1) to give 2-(6-amino-pyridin-3-ylmethyl)-3-methyl-succinic acid (0.5g, 37.7%).

(b) 2-(6-tert-Butoxycarbonylamino-pyridin-3-ylmethyl)-2-ethoxycarbonyl-3-methyl-succinic acid diethyl ester

A solution of 2-(6-amino-pyridin-3-ylmethyl)-3-methyl-succinic acid (300mg, 1.15 mmol) and NaH (30 mg, 1.25 mmol) in DMF (1 mL) was stirred at room temperature for 1 h under argon. A solution of (5-bromomethyl-pyridin-2-yl)-carbamic acid tert-butyl ester (330 mg, 1.15 mmol) in DMF (1 mL) was added and the reaction mixture was stirred for 24 h. The reaction was poured into H₂O and extracted with ethyl acetate. The organic layer was washed with water, dried and concentrated under reduced pressure. The product was purified by column chromatography (petroleumether/ethyl acetate, 4:1) to give 2-(6-tert-butoxycarbonylamino-pyridin-3-ylmethyl)-2-ethoxycarbonyl-3-methyl-succinic acid diethyl ester (220 mg, 40.9%).

(c) 2-(6-Amino-pyridin-3-ylmethyl)-3-methyl-succinic acid

A solution of 2-(6-tert-butoxycarbonylamino-pyridin-3-ylmethyl)-2-ethoxycarbonyl-3-methyl-succinic acid diethyl ester (94 mg, 0.20 mmol) in concentrated aqueous HCl (4 mL) was stirred at reflux overnight. The mixture was concentrated under reduced pressure and freeze dried to give the title compound as the HCl salt (50mg, 90.3%)

¹H NMR (400 MHz, D₂O): δ 1.22-1.35 (m, 3H), 2.77-3.08 (m, 4H), 6.99-7.05 (d, 1H), 7.70 (s, 1H), 7.84-7.91 (m, 1H).

MS (+) 239 (M+1)

Example 20

2-(6-Amino-pyridin-3-ylmethyl)-3-phenethyl-succinic acid

(a) 2-Ethoxycarbonyl-3-phenethyl-succinic acid diethyl ester

A solution of diethylmalonate (240.2 mg, 1.5 mmol) and NaH (43.2 mg, 1.8 mmol) in THF (3 ml) was stirred for one hour at room temperature under argon. 2-(4-Nitro-benzenesulfonyloxy)-4-phenyl-butyric acid ethyl ester (590 mg, 1.5 mmol) and DMPU (192.2 mg, 1.5 mmol) was added and the mixture was stirred at room temperature for three days. The reaction was poured into H₂O and extracted with ethyl acetate. The organic layer was washed with water and concentrated under reduced pressure. The product was purified

by column chromatography (petroleum ether/ethyl acetate, 4:1) to give 2-ethoxycarbonyl-3-phenethyl-succinic acid diethyl ester (274 mg, 52.1%).

(b) 2-(6-tert-Butoxycarbonylamino-pyridin-3-ylmethyl)-2-ethoxycarbonyl-3-phenethyl-succinic acid diethyl ester

A solution of 2-(6-amino-pyridin-3-ylmethyl)-3-phenethyl-succinic acid (9.3 mg, 0.027 mmol) and NaH (1.39 mg, 0.032 mmol) in DMF (0.5 ml) was stirred for 15 minutes at room temperature. (5-Bromomethyl-pyridin-2-yl)-carbamic acid tert-butyl ester (8.76 mg, 0.031 mmol) was added and the mixture was stirred at room temperature over night. The reaction was poured into H₂O and extracted with ethyl acetate. The organic layer was washed with water and concentrated under reduced pressure. The product was purified by column chromatography (heptane/ethyl acetate, 2:1) to give 2-(6-tert-butoxycarbonyl-amino-pyridin-3-ylmethyl)-2-ethoxycarbonyl-3-phenethyl-succinic acid diethyl ester (3.8 mg, 25.7 %).

(c) 2-(6-Amino-pyridin-3-ylmethyl)-3-phenethyl-succinic acid

A solution of 2-(6-tert-butoxycarbonylamino-pyridin-3-ylmethyl)-2-ethoxycarbonyl-3-phenethyl-succinic acid diethyl ester (94 mg, 0.169 mmol) in concentrated HCl (7 ml) was stirred at reflux for 24 hours. The mixture was concentrated under reduced pressure and freeze dried to give the title compound as the HCl salt (55 mg, 99.2 %).

¹H NMR (400 MHz, D₂O): δ 1.63-1.98 (m, 2H), 2.42-2.85 (m, 4H), 3.00-3.60(m, 2H), 6.85-6.98 (m, 1H), 7.10-7.29 (m, 5H), 7.76-8.05 (m, 2H).

MS (+) 329 (M+1)

Example 21

2-(6-Amino-pyridin-3-ylmethyl)-3-butyl-succinic acid

(a) 2-Butyl-3-ethoxycarbonyl-succinic acid diethyl ester

A solution of diethylmalonate (3.589 g, 0.022 mol) and cesium fluoride (3.406 g, 0.024 mol) in DMF (50 ml) was stirred for one hour at room temperature under argon. 2-Bromohexanoic acid ethyl ester (5 g, 0.022 mol) was added and the mixture was stirred at 100 °C and then at 65°C over night. The reaction was poured into H₂O and extracted with ethyl

acetate. The organic layer was washed with water and concentrated under reduced pressure. The product was purified by column chromatography (heptane/ethyl acetate, 1:2) to give 2-butyl-3-ethoxycarbonyl-succinic acid diethyl ester (5.0 g, 73.8 %).

5 (b) 2-(6-tert-Butoxycarbonylamino-pyridin-3-ylmethyl)-3-butyl-2-ethoxycarbonyl-succinic acid diethyl ester

A solution of 2-butyl-3-ethoxycarbonyl-succinic acid diethyl ester (1 g, 3 mmol) and NaH (119 mg, 5 mmol) in DMF (20 ml) was stirred for 60 minutes at 0°C under argon. (5-Bromomethyl-pyridin-2-yl)-carbamic acid tert-butyl ester (1.425 g, 5 mmol) was added
10 and the mixture was stirred at room temperature for one week. Ethanol (1 ml) was added and the reaction was poured into H₂O and extracted with ethyl acetate. The organic layer was washed with water and concentrated under reduced pressure. The product was purified by column chromatography (heptane/ethyl acetate, 4:1 to 1:1) to give 2-(6-tert-butoxycarbonylamino-pyridin-3-ylmethyl)-3-butyl-2-ethoxycarbonyl-succinic acid diethyl
15 ester (1.2 g, 71.3 %).

(c) 2-(6-Amino-pyridin-3-ylmethyl)-3-butyl-succinic acid

A solution of 2-(6-tert-butoxycarbonylamino-pyridin-3-ylmethyl)-3-butyl-2-ethoxycarbonyl-succinic acid diethyl ester (50 mg, 0.098 mmol) in concentrated HCl (4
20 ml) was stirred at reflux for 24 hours. The mixture was concentrated under reduced pressure and freeze dried to give the title compound as the HCl salt (28 mg, 89.9 %).

¹H NMR (400 MHz, D₂O): δ 0.83-0.91 (m, 3H), 1.18-1.40 (m, 4H), 1.50-1.82 (m, 2H), 2.67-2.75 (m, 12H), 2.78-2.85 (m, 2H), 2.89-3.03 (m, 1H), 6.97-7.09 (m, 1H), 7.63-7.67 (m, 1H), 7.80-7.85 (m, 1H).

25 MS (+) 281 (M+1)

Abbreviations

Ac = acetate

aq = aqueous

30 AIBN = α,α'-azoisobutyronitrile


Bn = benzyl

BSA = N,O-bis(trimethylsilyl)acetamide

- Bu = butyl
Bz = benzoyl
DCC = dicyclohexylcarbodiimide
DIAD = diisopropyl azodicarboxylate
5 DIPEA = diisopropylethylamine
DMAP = N,N-dimethyl amino pyridine
DME = 1,2-dimethoxyethane
DMF = dimethylformamide
DMSO = dimethylsulfoxide
10 EDC = 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide
Et = ethyl
EtOAc = ethyl acetate
EtOH = ethanol
h = hour
15 HMDS = hexamethyldisilazane
HOAc = acetic acid
HOBt = 1-hydroxybenzotriazol
HPLC = high performance liquid chromatography
KHMDs = potassium bis(trimethylsilyl)amide
20 LDA = lithium diisopropylamide
Me = methyl
MeOH = methanol
min = minutes
PMB = 4-methoxybenzyl
25 Ph = phenyl
Pr = propyl
PyBOP = (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate
TEA = triethylamine
TFA = trifluoroacetic acid
30 THF = tetrahydrofuran
TMSCN = trimethylsilyl cyanide
Tos = toluene-4-sulfonyl

CLAIMS

1. A compound of general Formula I



or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt, wherein

R₁ represents,

C₁-C₆ alkyl, substituted with one or more basic groups such as amino, amidino and/or guanidino;

cycloalkyl, substituted with one or more basic groups such as amino, amidino and/or guanidino;

heterocyclyl, containing at least one nitrogen atom;

heterocyclyl, containing at least one hetero atom selected from S or O,




and substituted with one or more basic groups such as amino, amidino and/or guanidino;

or aryl, substituted with one or more basic groups such as amino, amidino and/or guanidino,

R₂ represents H, acyl, acylamino, alkyl, alkylcarbamoyl, alkylthio, alkoxy, aroyl,

aroylamino, aryloxy, arylthio, amidino, amino, aryl, carbamoyl, carboxy, cyano, cycloalkyl, formyl, guanidino, halogen, heterocyclyl, hydroxy, oxo, nitro, thiol, Z₂N-CO-O-, ZO-CO-NZ- or Z₂N-CO-NZ- group,

R₃ represents COOR₅, SO(OR₅), SO₃R₅, P=O(OR₅)₂, B(OR₅)₂, P=OR₅(OR₅), or tetrazole, or any carboxylic acid isostere,

R₄ represents a  -group, or a  -group, or a  -group, roup,

R₅ represents H, C₁-C₆ alkyl or aryl,

R₆ represents C₁-C₆ alkyl, aryl, cycloalkyl, heterocyclyl, or an optionally N-substituted

H₂N-C(Z)-CONH-C(Z)- or H₂N-C(Z)- group,

5 R₇ represents H or C₁-C₆ alkyl,

X represents O, S, SO, SO₂, C(Z)₂, N(Z), NR₇SO₂, SO₂NR₇, NR₇CO or CONR₇,

Y represents O, N(Z), S, C(Z)₂, or a single bond,

Z represents independently H, C₁-C₆ alkyl, aryl, cycloalkyl or heterocyclyl,

with the proviso that when X represents O, S, SO, SO₂, N(Z), NR₇SO₂, SO₂NR₇, or

10 NR₇CO then Y represents C(Z)₂ or a single bond.

2. The compound according to claim 1, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt,

wherein

15 R₁ represents,

cycloalkyl, substituted with one or more basic groups such as amino, amidino and/or guanidino;


heterocyclyl, containing at least one nitrogen atom;

heterocyclyl, containing at least one hetero atom selected from S or O, and substituted

20 with one or more basic groups such as amino, amidino and/or guanidino;

R₂ represents H, C₁-C₃ alkyl, amino, halogen, hydroxy,

R₃ represents COOR₅,

R₄ represents a  -group,

25 R₅ represents H, C₁-C₆ alkyl or aryl,

R₆ represents C₁-C₆ alkyl, aryl, cycloalkyl, heterocyclyl, or an optionally N-substituted
H₂N-C(Z)-CONH-C(Z)- or H₂N-C(Z)- group,

X represents C(Z)₂,

Y represents O or C(Z)₂,

5 Z represents independently H or C₁-C₆ alkyl.

3. The compound according to claim 1, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt, wherein

10 R₁ represents,

cycloalkyl, substituted with one or more basic groups such as amino, amidino and/or guanidino;


heterocyclyl, containing at least one nitrogen atom;

heterocyclyl, containing at least one hetero atom selected from S or O, and substituted

15 with one or more basic groups such as amino, amidino and/or guanidino;

R₂ represents H, C₁-C₃ alkyl, amino, halogen or hydroxy,

R₃ represents COOR₅,

R₄ represents a  -group,

R₅ represents H, C₁-C₆ alkyl or aryl,

20 R₇ represents H or C₁-C₆ alkyl,

X represents C(Z)₂,

Y represents C(Z)₂, or a single bond,

Z represents independently H or C₁-C₆ alkyl.

25 4. The compound according to claim 1, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt, wherein

R₁ represents,

cycloalkyl, substituted with one or more basic groups such as amino, amidino and/or guanidino;


heterocyclyl, containing at least one nitrogen atom;

heterocyclyl, containing at least one hetero atom selected from S or O,

5 and substituted with one or more basic groups such as amino, amidino and/or guanidino;

R₂ represents H, C₁-C₃ alkyl, amino, halogen or hydroxy,

R₃ represents COOR₅,

10 R₄ represents a -R₅-group,

R₅ represents H, C₁-C₆ alkyl or aryl,


X represents C(Z)₂,

Y represents C(Z)₂, or a single bond,

Z represents independently H or C₁-C₆ alkyl.

15


5. A process for the preparation of a compound according to any one of claims 1-4, wherein R₁, R₅, R₆, and Z are as defined in claim 1 and R₂ is H, R₃ is COOR₅,

R₄ represents a -group,


X is C(Z)₂, Y is C(Z)₂,

comprising the step of:

20 reacting a compound of Formula VI,



wherein R_1 and Z is as defined in claim 1 and X is $\text{C}(\text{Z})_2$ and Y is $\text{C}(\text{Z})_2$ with a compound of Formula IX,




5

wherein R_6 is as defined in claim 1, in the presence of a suitable reagent, such as BSA or HMDS, under standard conditions.

6. A process for the preparation of a compound according to any one of claims 1-4,

10 wherein R_1 , R_2 , R_5 , R_6 , and Z are as defined in claim 1, R_3 is COOR_5 , X is $\text{C}(\text{Z})_2$, Y is O ,



and

comprising the step of:

reacting a compound of Formula XII,

15




wherein R_1 and R_2 are as defined in claim 1 and X is $\text{C}(\text{Z})_2$ with a compound of Formula XIII,



20

wherein R_6 is as defined in claim 1, in the presence of suitable coupling reagents such as DCC/DMAP, PyBop/DIPEA or SOCl_2 , under standard conditions.

7. A process for the preparation of a compound according to any one of claims 1-4,
 5 wherein R_1 and R_2 are as defined in claim 1 and X and Y is $\text{C}(\text{Z})_2$ or a single bond and R_3 and R_4 are COOR_5 , comprising the step of:
 reacting a compound of Formula XIV,




- 10 wherein R_2 and R_5 are as defined in claim 1 and Y is $\text{C}(\text{Z})_2$ or a single bond, with a
 compound of the general Formula III,




- wherein R_1 is as defined in claim 1, X is $\text{C}(\text{Z})_2$ and L is a suitable leaving group, such as Cl, Br, I or tosyl, in the presence of a suitable base, such as LDA or NaH, under standard
 15 conditions.

8. A process for the preparation of a compound according to any one of claims 1-4,
 wherein R_1 , R_2 , R_5 , R_7 , X, Y and Z are as defined in claim 1, R_3 is COOR_5 and



20

comprising the step of:
 reacting a compound of Formula XV,



with a compound of Formula XVI,



wherein R_7 is as defined in claim 1, in the presence of suitable reagents, such as DCC/DMAP, under standard conditions.

9. A pharmaceutical formulation containing a compound according to any one of claims 1-4 as active ingredient in combination with a pharmaceutically acceptable adjuvant, diluent or carrier.
10. The use of a compound according to any one of claims 1 to 4 in therapy.
11. The use of a compound according to any one of claims 1 to 4 for the manufacture of a medicament for the inhibition of carboxypeptidase U.
12. A method for treatment or prophylaxis of conditions associated with inhibition of carboxypeptidase U, comprising administering to a mammal, including man, in need of such treatment an effective amount of a compound as defined in any of claims 1-4.
13. A pharmaceutical formulation for use in the treatment or prophylaxis of conditions associated with inhibition of carboxypeptidase U, comprising a compound as defined in any one of claims 1-4 in combination with a pharmaceutically acceptable adjuvant, diluent or carrier.
14. A pharmaceutical formulation, comprising:

(i) a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt, and
(ii) one or more antithrombotic agent with a different mechanism of action, such as an antiplatelet agent, thromboxane receptor inhibitor, synthetase inhibitor, fibrinogen receptor antagonist, prostacyclin mimetic, phosphodiesterase inhibitor or ADP-receptor (P₂T) antagonist,
in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier.

15. A kit of parts comprising:

- 10 (i) a pharmaceutical formulation containing a compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier; and
(ii) a pharmaceutical formulation containing one or more antithrombotic agent with a different mechanism of action, such as an antiplatelet agent, thromboxane receptor inhibitor, synthetase inhibitor, fibrinogen receptor antagonist, prostacyclin mimetic,
15 phosphodiesterase inhibitor or ADP-receptor (P₂T) antagonist, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier;

which compound (i) and agent (ii) are each provided in a form that is suitable for administration in conjunction with the other.

20

16. A method for treatment of a patient suffering from, or susceptible to, a condition in which inhibition of carboxypeptidase U and a different antithrombotic mechanism are required or desired, which method comprises administering to the patient a therapeutically effective total amount of

- 25 (i) a compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier; in conjunction with
(ii) one or more antithrombotic agent with a different mechanism of action, such as an antiplatelet agent, thromboxane receptor inhibitor, synthetase inhibitor, fibrinogen
30 receptor antagonist, prostacyclin mimetic, phosphodiesterase inhibitor or ADP-receptor (P₂T) antagonist,

in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier.

17. A method for treatment of a patient suffering from, or susceptible to, a condition in which inhibition of carboxypeptidase U and a different antithrombotic mechanism are required or desired, which method comprises administering to the patient a formulation as defined in claim 15.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 00/00846

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07D 9/38, C07D 213/54, C07D 211/34, A61K 31/662, A61K 31/44, A61K 31/445
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

24 August 2000

Date of mailing of the international search report

28-08-2000

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Göran Karlsson/gh
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE00/00846

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 16-17
because they relate to subject matter not required to be searched by this Authority, namely:
A method for treatment of the human or animal body by therapy,
see rule 39.1
2. ☒ Claims Nos.: 1-15
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
see next sheet

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).:

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
pct/SE00/00846

The present claims 1-15 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. No special search effort can be made for searching unduly wide and speculative claims (PCT Search Guidelines C-III 3.7).

Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts related to the compounds prepared in the examples.

The applicants attention is drawn to the fact that claims relating to inventions in which no international search report has been established will not be the subject of an international preliminary examination (Rule 66.1(e) PCT). This is the case irrespective of whether or not the claims are amended following receipt of the search report during any Chapter II procedure.

REVISED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 November 2000 (09.11.2000)

PCT

(10) International Publication Number
WO 00/66550 A1

(51) International Patent Classification⁷: C07F 9/38, C07D
213/54, 211/34, A61K 31/662, 31/44, 31/445

(21) International Application Number: PCT/SE00/00846

(22) International Filing Date: 3 May 2000 (03.05.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9901572-9 3 May 1999 (03.05.1999) SE

(71) Applicant (for all designated States except US): AS-
TRAZENECA AB [SE/SE]; S-151 85 Södertälje (SE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): LINSCHOTEN,
Marcel [NL/SE]; AstraZeneca R & D Mölndal, S-431 83
Mölndal (SE). POLLA, Magnus [SE/SE]; AstraZeneca R
& D Mölndal, S-431 83 Mölndal (SE).

(74) Agent: ASTRAZENECA AB; Global Intellectual Prop-
erty, Patents, S-151 85 Södertälje (SE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ,
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,
TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:


— With international search report.

(88) Date of publication of the revised international search
report: 25 January 2001

(15) Information about Correction:
see PCT Gazette No. 04/2001 of 25 January 2001, Section
II

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: NEW COMPOUNDS



(57) Abstract: The present invention relates to compounds of Formula (I), and pharmaceutically acceptable salts or solvates thereof, or solvates of such salts, which compounds inhibit carboxypeptidase U and thus can be used in the prevention and treatment of diseases associated with carboxypeptidase U. In further aspects, the invention relates to compounds of the invention for use in therapy; to processes for preparation of such new compounds; to pharmaceutical compositions containing at least one compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, as active ingredient; and to the use of the active compounds in the manufacture of medicaments for the medical use indicated above.

WO 00/66550 A1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 00/00846

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07F 9/38, C07D 213/54, C07D 211/34, A61K 31/662, A61K 31/44, A61K 31/445
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

24 August 2000

Date of mailing of the international search report

12 6 -10- 2000

Name and mailing address of the ISA/

Swedish Patent Office

Authorized officer

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE00/00846

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 16-17
because they relate to subject matter not required to be searched by this Authority, namely:
**A method for treatment of the human or animal body by therapy,
see rule 39.1**
2. ☒ Claims Nos.: 1-15
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
see next sheet
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a):

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

pct/SE00/00846

The present claims 1-15 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. No special search effort can be made for searching unduly wide and speculative claims (PCT Search Guidelines C-III 3.7).

Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts related to the compounds prepared in the examples.

The applicants attention is drawn to the fact that claims relating to inventions in which no international search report has been established will not be the subject of an international preliminary examination (Rule 66.1(e) PCT). This is the case irrespective of whether or not the claims are amended following receipt of the search report during any Chapter II procedure.